

Terminology and Ontology Development for Semantic Annotation: A Use Case on Sepsis and Adverse Events

Melissa Y. Yan ^{a,*}, Lise Tuset Gustad ^{b,c,d}, Lise Husby Høvik ^{c,e} and Øystein Nytrø ^a

^a *Department of Computer Science, Norwegian University of Science and Technology, Trondheim, Norway*

E-mails: melissa.yan@ntnu.no, nytroe@ntnu.no

^b *Faculty of Nursing and Health Sciences, Nord University, Levanger, Norway*

E-mail: lise.t.gustad@nord.no

^c *Department of Circulation and Medical Imaging, Norwegian University of Science and Technology, Trondheim, Norway*

E-mails: lise.t.gustad@ntnu.no, lise.hovik@ntnu.no

^d *Department of Medicine, Levanger Hospital, Nord-Trøndelag Hospital Trust, Levanger, Norway*

E-mail: lise.gustad@hnt.no

^e *Clinic of Anaesthesia and Intensive Care, St. Olavs hospital, Trondheim University Hospital, Trondheim, Norway*

E-mail: lise.husby.hovik@stolav.no

Abstract. Annotations enrich text corpora and provide necessary labels for natural language processing studies. To reason and infer underlying implicit knowledge captured by labels, an ontology is needed to provide a semantically annotated corpus with structured domain knowledge. Utilizing a corpus of adverse event documents annotated for sepsis-related signs and symptoms as a use case, this paper details how a terminology and corresponding ontology were developed. The Annotated Adverse Event NOte TErminology (AAENOTE) represents annotated documents and assists annotators in annotating text. In contrast, the complementary Catheter Infection Indications Ontology (CIIO) is intended for clinician use and captures domain knowledge needed to reason and infer implicit information from data. The approach taken makes ontology development understandable and accessible to domain experts without formal ontology training.

Keywords: ontology development, clinical knowledge representation and reasoning, semantic annotation, sepsis, adverse events

1. Introduction

Many natural language processing (NLP) studies rely on annotated corpora to create models for text classification, information extraction, named entity recognition, question answering, summarization, and text generation. Often, semantic annotation is done to capture domain knowledge within the text. Annotated corpora are frequently generated by annotators based on an annotation guideline, which provides the stan-

dard and rules for how to label text using specified terms. This annotation guideline is usually similar to a terminology unless the corpus was annotated using an ontology. To further enrich an annotated corpus by capturing, reasoning, and inferring the underlying associated domain knowledge, an ontology is needed.

To demonstrate semantic annotation terminology and ontology development, the use case is based on clinician-presented needs for identifying sepsis from adverse events (AEs). Improperly cared for peripheral intravenous catheter (PIVC) medical devices can lead to unwanted and unintentional events that harm pa-

*Corresponding author. E-mail: melissa.yan@ntnu.no.

tients, such as AEs like phlebitis, bloodstream infections (BSIs), and sepsis. However, PIVCs are poorly documented in clinical records because of routine use among inpatients, and sepsis is also poorly documented outside of the intensive care unit (ICU). The lack of explicitly documented concepts makes it challenging to directly detect and annotate PIVC-related phlebitis, BSI, and sepsis for quality surveillance to improve care. Thus, indications for the presence of PIVCs, phlebitis, infections, and sepsis are annotated instead. Those annotations are structurally preserved as a terminology and the clinical knowledge required to reason about the indications is represented in an ontology. Additional details for the use case are provided in Section 2.

This paper provides a detailed and concrete description of the methodology utilized for ontology development of an annotated corpus based on a use case from the clinical domain. The main contributions presented are:

1. Describing the development process for constructing a terminology that can represent an annotated corpus. Specifically, a terminology for indexing annotated AE documents.
2. Presenting the development process for the terminology's corresponding ontology, which represents domain knowledge and allows inferring implicit knowledge in a specific domain. The corresponding ontology in the use case represents clinical domain knowledge specifically for annotated catheter-related and infection-related signs in AE documents.
3. Releasing a terminology and ontology that can be applicable to identifying and reasoning about sepsis in an AE corpus.

This paper significantly extends the papers [1] and [2], by adding an ontology with instances and including evaluation of the correctness and ability to answer competency questions. In addition, instances from the annotated corpus in [2] were added into the terminology, the terminology was evaluated using competency questions, and an ontology was developed to answer competency questions with clinical knowledge.

Based on the presented use case in Section 2 and objective in Section 2.2, an Annotated Adverse Event Note Terminology (AAENOTE) and corresponding Catheter Infection Indications Ontology (CIIO) were developed. Section 5 details the terminology construction and development process to represent annotated documents, and Section 6 presents the results and eval-

uations for the AAENOTE. To address shortcomings in the terminology for annotated documents, Section 7 describes the ontology development process for domain knowledge representation of documented content. Additionally, Section 8 presents the results and evaluations for the CIIO. Finally, Section 9 discusses the findings, limitations, representations, accessibility, and utility.

2. Use Case Background and Motivation

2.1. Sepsis from Peripheral Intravenous Catheter-related Phlebitis and Infection Adverse Events

As the most commonly used medical device in hospitals, PIVCs, are inserted into the peripheral vein to administer intravenous (IV) fluids, IV medications, and blood transfusions [3]. Improper management of PIVC or the infusions connected to the PIVC can lead to phlebitis, which is either infectious, mechanical, or chemical inflammation of the vein [4–6]. Independent of cause, all PIVC phlebitis share many symptoms like redness and swelling near a patient's infusion insertion site across infectious, mechanical, or chemical phlebitis, making it difficult to distinguish. Further, all PIVC-related phlebitis causes AEs like significant pain, PIVC failure that delays treatment, and compromising future venous access. Infectious phlebitis may lead to BSI due to: 1. migration of bacteria at the insertion site, 2. bacteria migrating through the catheter tract or catheter hub, 3. contaminated infusate, or 4. bacteria from an existing infection in the bloodstream attaching to the catheter [7]. BSIs can potentially to cause sepsis and occur when bacteria enter the bloodstream [8]. *Staphylococcus aureus* (*S. aureus*) is a lethal bacteria frequently found on skin that commonly causes BSIs [9], defined as a dysregulated host immune response to infection that results in organ failure and a mortality rate of 20% [10]. Approximately 7.6% to 35% of *S. aureus* BSIs are caused by PIVCs [11].

Even though PIVCs are frequently used, they are routinely not documented in clinical records [3]. Moreover, sepsis is also poorly documented outside the ICUs [12]. This lack of documentation makes it challenging to perform retrospective and real-time systematic quality surveillance of PIVC-related phlebitis or BSIs to identify learning opportunities for improving PIVC care to lower phlebitis, BSI-related, and PIVC-related AE incidents. Hence, AE reports or documents,

1 which are customarily used to report PIVC failures,
 2 were selected as the clinical text for this project. To
 3 capture documented observable patient states and infer
 4 underlying knowledge of PIVC-related phlebitis or
 5 BSI from clinical text, an ontology that models clinical
 6 knowledge representation and reasoning is necessary.

7 2.2. Use Case Objective

8
 9
 10 The use case objective was to develop a model for
 11 representing and reasoning about PIVC-related BSIs
 12 in the unstructured free-text of AE reports, describe
 13 the development process, and discuss the discoveries
 14 and limitations. From the research question “is there
 15 a connection between BSIs and PIVCs at the hospital?”,
 16 competency question requirements for an ontology
 17 representing and reasoning about PIVC-related
 18 BSIs were identified by clinicians as follows:

- 19 1. Does patientA have phlebitis, and was it infec-
- 20 2. Does patientA have an infection?
- 21 3. Does patientA have a BSI?
- 22 4. How many patients have an infection or BSI?
- 23 5. Which patients have sepsis?
- 24 6. Does patientB have a catheter?
- 25 7. Does patientB have a PIVC?
- 26 8. How many catheters does patientB have, where
- 27 9. Does patientC have an infection and catheter?
- 28 If so, was patientC’s infection associated with a
- 29 catheter?
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51

3. Related Work

3.1. Annotation, Tagging, and Ontologies for Natural Language Processing

40 Many studies focus on the relationship between an-
 41 notation, tagging, and ontologies for NLP develop-
 42 ment. These studies include annotating corpora, NLP
 43 extraction, and classification tasks. Below are some
 44 studies that have shared their findings, issues, and pos-
 45 sible solutions.

46 Annotated using the Uberon multi-species anatomi-
 47 cal ontology [13], the Colorado Richly Annotated Full-
 48 Text (CRAFT) Corpus is a resource for NLP develop-
 49 ment which is also semantically annotated with con-
 50 cepts from eight Open Biomedical Ontologies (OBO)
 51 and terminologies [14, 15]. However, while annotat-

1 ing with the OBOs, they discovered that the OBOs
 2 are not developed for annotation because there are
 3 overlapping terms within the different OBOs, context-
 4 specific definitions, and semantic ambiguities. Addi-
 5 tionally, some OBOs do not follow the OBO Foundry
 6 principle of using relations from the OBO Relation
 7 Ontology (RO) [16] to link concepts [17]. Therefore, to
 8 improve OBOs for semantic annotation of biomedical
 9 documents, the researchers proposed desirable ontol-
 10 ogy implementations such as, but not limited to, inte-
 11 grating overlapping OBOs terms, resolving ontology-
 12 specific ambiguities, and expanding relations [17].

13 A study comparing how anatomy ontologies are
 14 used for annotations discovered annotation and ontol-
 15 ogy issues [18]. Annotations from three public
 16 datasets were compared to anatomical terms in the
 17 Foundational Model of Anatomy (FMA) [19, 20] and
 18 Uberon [13] ontologies using the Zooma¹ and On-
 19 tology Mapper² software tools. Manual and semi-
 20 automated preprocessing were done to normalize
 21 terms, but there were few matches between the ontol-
 22 ogy and annotations, mainly because of strict match-
 23 ing. Additionally, the user-provided annotation labels
 24 resulted in mismatches, such as annotating a phrase
 25 with multiple ontology terms or using an abbrevia-
 26 tion or adjective for an anatomical part instead of the
 27 anatomical ontology term. Ontology issues include
 28 missing anatomical synonyms used by the annotators
 29 and differing anatomical terms in the ontologies be-
 30 cause the ontologies are designed for different pur-
 31 poses and made by different design decisions. The
 32 study concluded that mapping terms to an ontology re-
 33 quire a large amount of time, effort, and manual cura-
 34 tion. Further, an ontology’s design decisions and scope
 35 will affect users trying to match annotations to an on-
 36 tology, and ontologies must be used to understand their
 37 potential.

38 The Unified Medical Language System (UMLS) is
 39 a collection of standard biomedical terminology [21],
 40 and it has been used to process text by extracting
 41 concepts, relations, and knowledge (i.e., link or an-
 42 notate text with standard terminology) [22]. A soft-
 43 ware capable of finding and linking biomedical text to
 44 terminology concepts in the UMLS Metathesaurus is
 45 MetaMap [23]. However, the developers of MetaMap
 46 mention that improvement is required for detecting
 47 similar names, acronyms, and abbreviations and re-

¹<https://www.ebi.ac.uk/spot/zooma/>

²<http://www.ebi.ac.uk/efo/tools>

1 solving ambiguities by possibly distinguishing concepts using word sense disambiguation.

2 In [24], an overview of studies using knowledge bases for entity coreference resolution were discussed. Among those studies was the OntoNotes project, which annotated a multilingual corpus for different levels of semantic structure in the text [25–27]. One of the annotation levels includes linking OntoNotes word senses to the Omega ontology [27–29]. Near-synonymous word sense pools were created by specialists who grouped sense distinctions from WordNet and dictionaries based on similar definitions. This enables machines to automatically tag senses more accurately and improves inter-annotator agreement due to difficulties determining WordNet distinctions directly in the text [29]. Before each sense pool was linked to a concept in the Omega ontology [30], each sense pool was verified by machine and humans [29].

3.2. Ontology Development Methods and Evaluation

22 There are many ontology development methods, such as: the Enterprise ontology’s Uschold and King [31], the Toronto Virtual Enterprise (TOVE) ontology’s Grüninger and Fox [32], METHONTOLOGY [33], the On-To-Knowledge Methodology (OTKM) [34], and NeOn [35]. Of those methods, Uschold and King [31] and Grüninger and Fox [32] follow a sequential sequence of phases, whereas METHONTOLOGY [33], OTKM [34], and NeOn [35] are iterative. Whether sequential or iterative, the previously mentioned methods and 2 reviews [36, 37] have shown that ontology development typically includes the phases: specification, conceptualization, formalization, implementation, and maintenance. During those phases, the knowledge acquisition, evaluation, and documentation phases also commonly occur either as a separate phase or concurrently with other phases. Appendix A provides a summary of the methods for each phase.

40 During and between phases, ontology evaluation judges an ontology’s content to a reference, such as requirement specifications, competency questions [32], or the real-world [38, 39]. Evaluation includes: (1) verification that the ontology has the correct informal natural language definition and formal ontology language definition, and (2) validation that the ontology represents the world it was created for [38, 39]. In theory, there are many criteria for evaluation, but in practice, most studies only use the expressiveness and practical usefulness criteria [40]. Expressiveness is the number of competency questions answerable by the ontol-

ogy [32, 40, 41], and practical usefulness is the number of problems an ontology can be applied to [40, 41].

3.3. Relevant Ontology Resources

6 There lacks an ontology specifically for sepsis-related BSI, infection signs, anatomical locations, medical devices, and procedures. However, pre-existing ontologies can contain relevant concepts. For example, the Infectious Disease Ontology (IDO) [42] has sepsis and hospital-acquired infection entities. Sign and symptom entities are present in the Ontology for General Medical Science (OGMS) [43], and vital sign entities exist in the Vital Sign Ontology (VSO) [44, 45]. Anatomical locations can be described using anatomical entities of the Foundational Model of Anatomy Ontology (FMA) [19, 46] and anatomical spatial location descriptor entities from the Biological Spatial Ontology (BSPO) [47]. Because AE reports are used, the adverse event entities in the Ontology of Adverse Events (OAE) [48, 49] might also be relevant. Furthermore, relationship object properties in the Open Biological and Biomedical Ontology (OBO) Relation Ontology [16, 50] could be used to link different entities together to capture more information.

26 In addition to ontologies, there are also potential relevant terminologies and taxonomies. For example, there are different procedure, medical device, and catheter terms in the National Cancer Institute Thesaurus (NCIT) [51, 52]. Potential relevant standardized nursing practice language is found in the International Classification for Nursing Practice (ICNP) terminology [53], Nursing Interventions Classification (NIC) taxonomy [54], and NANDA International Nursing Diagnoses Classification taxonomy [55]. Furthermore, infusion phlebitis-related information can be obtained from the 1998 Visual Infusion Phlebitis Scale [56] and the 2021 Infusion Therapy Standards of Practice Updates [57]. Concepts or terms from these resources can be used to expand the ontology if deemed necessary by ontology users. The relevant ontologies, terminologies, taxonomies, and clinical guidelines can be found in Table 1.

4. Materials

4.1. Synthetic Dataset

50 Documents for annotation are from an AE synthetic dataset. The documents are based on unstruc-

Table 1
Overview of Relevant Resources for This Study

Type	Resource	Relevant Concepts or Terms
Ontology	Infectious Disease Ontology (IDO)	Sepsis and hospital-acquired infection entities
	Ontology for General Medical Science (OGMS)	Sign and symptom entities
	Vital Sign Ontology (VSO)	Vital sign entities
	Foundational Model of Anatomy Ontology (FMA)	Anatomical entities
	Biological Spatial Ontology (BSPO)	Anatomical spatial location descriptor entities
	Ontology of Adverse Events (OAE)	Adverse event entities
	Open Biological and Biomedical Ontology (OBO) Relation Ontology	Relationship object properties
Terminology	National Cancer Institute Thesaurus (NCIT) terminology	Procedure and medical device terms
	International Classification for Nursing Practice (ICNP) terminology	Terms
Taxonomy	Nursing Interventions Classification (NIC) taxonomy	Terms
	NANDA International Nursing Diagnoses Classification taxonomy	Terms
Clinical Guideline	1998 Visual Infusion Phlebitis Scale	Visual infusion phlebitis grading scale
	2021 Infusion Therapy Standards of Practice Updates	Updated infusion therapy practice standards




Fig. 1. From unannotated documents to an annotated corpus for populating instances in a terminology. A 100 synthetic adverse event (AE) unstructured free-text documents were manually generated. Those synthetic documents were annotated by 8 annotators over 4 annotation sessions using revised annotation guidelines. Each annotator annotated 70 documents (i.e., 10 documents in session 1 and 20 documents in the remaining 3 sessions) to produce a total of 560 annotated synthetic AE documents.

ured free-text AE notes within the extracted AE reports from the electronic incident reporting system at St. Olavs hospital, Trondheim University Hospital in Trondheim, Norway, between September 2015 to December 2019 [2]. The synthetic dataset contains 100 AE notes or documents manually created and verified by a nurse to ensure clinical data is anonymized. The Norwegian Regional Committees for Medical and Health Research Ethics (REK) has granted ethical approval to use AEs in this paper (approval no 2018/1201/REKmidt, 26814).

4.2. Annotated Synthetic Dataset

The synthetic documents were annotated by 8 annotators with clinical backgrounds over 4 annotation sessions [2]. Each annotator annotated 10 documents in session 1 and 20 documents in the remaining 3 sessions (i.e., 70 documents annotated over 4 annotation session). This resulted in 560 annotated synthetic AE documents as shown in Fig. 1 (i.e., 8 annotators * 70 annotated documents * 4 sessions = 560 total an-

notated synthetic AE documents). In each annotation session, annotators followed the annotation guideline and used the Brat rapid annotation tool (BRAT) [58] to annotate the documents. Then, documents were evaluated and manually screened to identify ambiguities for revising the annotation guideline. This process was repeated 3 additional times with a new set of documents and a revised annotation guideline.

4.3. Annotation Guideline

An annotation guideline was developed based on the clinical research question: Is there a connection between BSIs and PIVCs at the hospital? Discussions with nurses provided insight into how catheters can be distinguished explicitly by the name or implicitly based on the anatomical insertion site or procedure mentioned. This formed into four domain-specific questions of interest:

1. What are the different signs of infections, specifically for BSIs, sepsis, or infected PIVCs?

2. What are the signs for different types of catheters?
3. Where are the anatomical insertion sites of catheters?
4. What procedures, interventions, and activities can be related to catheter use?

Answers to domain-specific questions were then sorted into the following 7 main categories:

1. **Sign**: infection signs
2. **Location**: anatomical insertion sites
3. **Device**: signs of catheter types
4. **Procedure**: procedures, interventions, or activities related to catheters
5. **Sensitivity**: protected health information
6. **Person**: individuals mentioned, such as patient, clinician, or relative.
7. **Whole**: label representing the span of the whole text and given to indicate if the document contains information about infection, BSI, sepsis, faulty device malfunctioning, catheter, PIVC, or is sensitive.

All categories except **Whole** have a hierarchy with more specific subcategories to capture detailed granularity from text (e.g., the **Person** category has a **Patient** subcategory). Furthermore, to capture relationships between categories for downstream analysis (e.g., infection sign at a specific location), the following four relationships to link categories were included:

1. **Person** $\xrightarrow{\text{Person has}}$ **Sign, Location, Device, or Procedure**
2. **Procedure** $\xrightarrow{\text{Procedure uses}}$ **Device**
3. **Sign** $\xrightarrow{\text{Caused by}}$ **Device or Procedure**
4. **Sign, Device, or Procedure** $\xrightarrow{\text{Located nearby/on/in}}$ **Location**

A preliminary annotation guideline was created using the seven categories and four relationships. Annotation guidelines from each of the 4 annotation sessions are available online³ [2].

5. Terminology Development for Annotations

In annotations, categories are known as entities for labeling a span of words or phrases. Whereas in ontologies and terminologies, categories are known as classes. To separate the annotation guideline from the terminology and ontology, the annotation categories and entities are in **bold font**, and the terminology and ontology classes are in `typewriter` font.

³<https://folk.ntnu.no/melissay/ae-guidelines/>

Table 2
Convert Annotation Guideline to Terminology

Annotation Guideline	Terminology
Entity (category) hierarchy	Class hierarchy
Attributes which provide detailed entity information	Data properties
Relationships between entities	Object properties

5.1. Design Decision for Annotations

The terminology was developed using the bottom-up approach based on the annotation guideline refinement process from 4 iterations. Competency questions were not used to create this terminology. This terminology is meant to assist annotators who want to label text and allow users interested in performing downstream analyses to adjust the granularity of labels. The objective is solely to represent the annotated corpus and provide structure to the terminology used by annotators. Thus, included individuals are based on concrete examples from the annotated corpus. A simplified example of how annotation labels in annotated documents are added to the terminology as individuals is provided in Fig. 2.

Instead of reusing and re-defining existing ontologies, it was easier and simpler to develop a terminology based on what is documented in the data. For instance, although the FMA contains relevant anatomical parts, the ontology was too complex and detailed to be incorporated easily into the terminology to fit the use case's purpose. Additionally, the purpose was to include only concrete items documented in the terminology and not provide terminology for all existing items. By opting to simplify the terminology, it was easier to build the terminology directly based on the annotation guideline and modify the terminology to incorporate feedback from discussions with clinicians.

5.2. Convert Annotation Guideline To Terminology

The categories, attributes, and relationships in annotation guidelines described in Section 4.3 correspond to classes, data properties, and object properties in terminologies (Table 2 and Fig. 3).

The terminology was developed from the annotation guideline by translating each hierarchy of entities into a class hierarchy, using attribute information to add data properties, and converting relationships into object properties. During development, the terminology was modified to remove ambiguities by adding new




Fig. 2. Using an annotated corpus to populate individuals in a terminology. Each of the 560 documents were translated into an individual and each label within a document was also translated into an individual. In the simplified example, an annotated document has 2 **patient** labels, 2 **PIVC** labels, and 2 $\xrightarrow{\text{Person has}}$ relationships linking the labels. Each label is converted into an individual (i.e., a purple diamond) of the corresponding class (i.e., **Patient** or **PIVC** yellow circle). Then the labels are linked using the $\xrightarrow{\text{Person has}}$ object property, similarly to how the $\xrightarrow{\text{Person has}}$ relationship links labels in the annotated text.




Fig. 3. Terminology development. The annotation guideline from the fourth annotation session was converted into a terminology. Annotation categories were converted into ontology classes, relationships into object properties, and attributes into data properties. Then the individuals of documents and labels were added. Additional modifications were incorporated as needed, such as removing ambiguities, re-organizing hierarchies, and adding missing concepts. This resulted in the AAENOTE which models and provides an index of annotated documents.

class hierarchies and modifying class names, object properties, and data properties. Specifically, to remove ambiguity between symptoms, infection signs, and device malfunction signs, the **Observation** class was introduced to encompass them. Additionally, the **Sensitive** category was revised to the **Identifier** class and the $\xrightarrow{\text{Caused by}}$ relationship was revised to the $\xrightarrow{\text{Is observed with}}$ relationship. A summary of the converted 7 main classes can be found in Table 3.

The results from all 4 annotation sessions were included as individuals in the terminology, but the terminology only reflects results based on the last annotation guideline. To accommodate revisions in the annotation guideline, annotation categories that were revised in the guideline are updated in the terminology correspondingly. For instance, removed annotation categories are reflected by changing the granularity of the removed category to a higher level. Annotation categories can also be re-organized to become subclasses of a different class. Moreover, newly added annotation categories are directly added as new classes in the terminology. An example is depicted in Fig. 4.

Although the terminology was not developed to answer competency questions, the competency questions were still used to determine what could be found in an-

notated documents. To answer competency questions, the annotated documents were imported into the terminology as individuals.

6. Terminology Results for Annotations

6.1. Annotated Adverse Event NOte Terminology (AAENOTE)

Annotated AE documents and their annotations are modeled by the Annotated Adverse Event NOte Terminology (AAENOTE). To increase accessibility, the terminology is in both English and Norwegian. There are 149 classes, 5 object properties, 27 data properties, and 4470 individuals. The 7 classes which form the main hierarchies are:

1. **Observation**: Any sign or symptom that can be monitored.
2. **Anatomical Location**: Any anatomical body part, organ, or relative position of the body.
3. **Medical Device**: Any instrument, device, or equipment used for a medical purpose.
4. **Procedure**: Any procedure, intervention, or activity related to catheters.

Table 3

Annotation Guideline vs Annotated Adverse Event NOTE Terminology (AAENOTE). Annotation categories and entities are in **bold font** and the terminology and ontology classes are in typewriter font

Annotation Categories	Category Description	Sub-categories	Terminology Classes	Class Description	Subclasses
Sign	Infection signs	29	Observation	Documented clinical observation including symptoms, infection signs, and device malfunctions	41
Location	Anatomical insertion sites	17	Anatomical Location	Anatomical location	25
Device	Signs of catheter types	16	Medical Device	Treatment equipment or part	19
Procedure	Procedures, interventions, or activities related to catheters	31	Procedure	Procedure, intervention, or activity for catheter-related versus non-catheter related	36
Sensitivity	Protected health information	14	Identifier	Protected health information	14
Person	Individuals	3	Person	Individual	3
Whole	Label representing whole text indicating the note contains infection, BSI, sepsis, faulty device malfunctioning, catheter, PIVC, or sensitive information	0	Annotated Document	Representation of an AE note's filename, annotation session, annotator, and annotated labels	4




Fig. 4. Handling annotation guideline revisions in the terminology. If the **Delete Device** annotation category in red is removed from the annotation guideline, then the **Delete Device** class is removed from the terminology and all individuals of the **Delete Device** terminology class are now part of the superclass **Medical Device**. If the **PIVC** (peripheral intravenous catheter) annotation category in orange is re-organized to become the sub-category of **Venous Catheter**, then the **PIVC** class is now a subclass of **Venous Catheter** and the individuals remain as part of the **PIVC** class. If the **Arterial Catheter** annotation category in gray is added, then the **Arterial Catheter** class is added to the terminology and the corresponding individuals will be added as well.

- 6. **Person**: An individual, such as a patient, clinician, or relative.
- 7. **Annotated Document**: Annotated adverse event document metadata and labels.

Relationships between the 7 class hierarchies can be formed using the following 5 object properties:

1. **Person** $\xrightarrow{\text{Person has}}$ **Observation, Anatomical Location, Medical Device, or Procedure**
2. **Procedure** $\xrightarrow{\text{Procedure uses}}$ **Medical Device**
3. **Observation** $\xrightarrow{\text{Is observed with}}$ **Medical Device or Procedure**
4. **Observation, Medical Device, or Procedure** $\xrightarrow{\text{Located nearby/on/at/in}}$ **Anatomical Location**
5. **Annotated Document** $\xrightarrow{\text{Has label}}$ **Anatomical Location, Identifier, Medical Device, Observation, Person, or Procedure**

An example showing AAENOTE representing an annotated document using parts of the class hierarchies and class properties is shown in Fig. 5. The complete class hierarchies of AAENOTE are in Appendix B.1.

Each annotated AE note or document is an individual of the **Annotated Document** class and can have object properties linking the AE document to individual labels from the other 6 class hierarchies. Additionally, each AE document and has data properties for the filename, annotation session, and annotator. The individuals of the other 6 hierarchies can also have object properties and data properties if an annotator provides that information.

- 5. **Identifier**: Protected health information that can be used to identify an individual.




Fig. 5. Annotated Adverse Event Note Terminology (AAENOTE) Representing an Annotated Document. (a) Example of an annotated document with annotation categories and relationships that link the categories together. (b) Part of the terminology class hierarchies used within the annotation example are shown in the white boxes. The 3 annotated relationships (i.e., $\frac{\text{Person has}}{\text{Located nearby/on/at/in}}$, and $\frac{\text{Procedure uses}}{\text{}}$) are represented by the 3 object properties that link the classes together. Object properties are shown using the thicker colored lines with arrows.

6.2. AAENOTE Evaluation

The purpose of AAENOTE is to model and provide semantic meaning to annotated AE notes or documents. The terminology was not developed based on competency questions, but it would be interesting to see what competency questions could be answered. Hence, AAENOTE was evaluated using the competency questions as requirements. Competency questions using AAENOTE can only be answered based on explicitly annotated classes or subclasses. Words or phrases that lack annotation are excluded from this terminology. Thus, only the annotation category labels provided by annotators are included in the individuals as classes.

Knowledge represented by the terminology can either be found explicitly, based on the direct classes and relationships, or be inferred implicitly, based on underlying concrete knowledge and indirect classes and relationships. For example, the competency question “Does patientA have an infection?” can be answered explicitly by finding an individual of the patient class who has an infection (i.e., $\text{Patient} \xrightarrow{\text{Person has}} \text{infection}$). It can also be answered implicitly by finding an anatomical location that has an infection (i.e., $\text{Infection} \xrightarrow{\text{Located nearby/on/at/in}} \text{anatomical location}$) because an anatomical location, in this

terminology, must be part of a person. However, if the infection is not explicitly mentioned, this terminology cannot implicitly determine what other observations combined indicate an infection.

SPARQL query results vary depending on the clinician’s interest in knowing how many instances a patient has for one class or a combination of that one class with other classes. For example, to find how many patients explicitly have an infection, the query can be written to find all instances where either: 1. individuals of the patient class have the object property “person has” to an individual of the infection class (i.e., $\text{Patient} \xrightarrow{\text{Person has}} \text{infection}$), or 2. the individuals of the patient class have the object property “person has” to an individual of the infection class and/or an individual of the observation class or its subclass (i.e., $\text{Patient} \xrightarrow{\text{Person has}} \text{infection}$ and other observation(s)). The different number of instances is shown in Table 4. The first query has 14 instances where a patient has an infection regardless of other observations. In contrast, the second query divides the 14 instances into Query Result 2-14 to show the number of instances where a patient has an infection with different combinations of other observations. To implicitly find patients that have an infection, the infection must be located at an anatomical location of a per-

Table 4
AAENOTE SPARQL Query Result: Patient has Infection

Query Result	Instances	# Observations	Person		Observation												
			patient		blood pressure	body temp	c reactive protein	consciousness level	infection	neurological and physiological	observation	pulse	red	respiratory rate	sepsis	swollen	
1	14	1	✓						✓								
2	1	1	✓						✓								
3	1	2	✓					✓	✓								
4	1	2	✓						✓				✓				
5	1	2	✓						✓					✓			
6	1	3	✓			✓	✓		✓								
7	1	3	✓			✓			✓					✓			
8	1	3	✓					✓	✓	✓							
9	1	3	✓					✓	✓		✓						
10	2	3	✓						✓				✓				✓
11	1	4	✓					✓	✓	✓		✓					
12	1	5	✓			✓	✓	✓	✓				✓				
13	1	6	✓			✓	✓	✓	✓				✓			✓	
14	1	8	✓			✓	✓	✓	✓	✓	✓	✓	✓	✓			

Query result 1: Patient $\xrightarrow{\text{Person has}}$ infection.

Query result 2-14: Patient $\xrightarrow{\text{Person has}}$ infection and other observation(s).

son. Queries about infections at an anatomical location can be written using Infection $\xrightarrow{\text{Located nearby/on/at/in}}$ anatomical location) or using Infection and other observation(s) $\xrightarrow{\text{Located nearby/on/at/in}}$ anatomical location if the clinician is curious about additional observations that were documented with the infection. In AAENOTE, there is only 1 instance where an anatomical location (i.e., skin) has an infection, as shown in Query Result 1 of Table 5. However, it is more informative for clinicians to look at additional potential observations that can indicate infection around a certain location; Table 5 Query Results 2-10 provide some other observations that were found located with skin. The clinical knowledge required to find indications of infections and catheters is lacking in AAENOTE.

Explicit queries and basic implicit queries in AAENOTE can answer 7 of the 9 competency questions. However, these queries still lack the clinical knowledge needed to include more implicit queries by combining additional observations, anatomical locations, and/or procedures to identify indications. In Appendix B, Table 7 provides the terminology classes and relationships used to form explicit and implicit

queries to explicitly and implicitly answer each competency question. Additionally, concrete underlying knowledge used to make inferences is also provided. Results can be found in Appendix B.

7. Ontology Development for Domain Knowledge

7.1. Design Decision for Domain Knowledge

In this use case, the clinicians' need is to focus on identifying and inferring a patient's state based on documented observations in AE documents related to PIVCs and BSIs. A patient's underlying state can be measured by monitoring devices that measure vital signs (e.g., blood pressure, pulse, and respiratory rate) or exhibited by observable signs and symptoms (e.g., pain, fever, chills, and mobility impairment). Those measurable and observable signs and symptoms are then documented by clinicians in the electronic health record (EHR) to record patient conditions and communicate with other clinicians. When an AE incident could have or has happened, clinicians will go through the documentation to recall what occurred and report it in a separate AE document.

Table 5
AAENOTE SPARQL Results for Observations Located at the Skin

Query Result	Instances	Observations					
		edema	infection	pain	red	swollen	warm
1	1		✓				
2	3			✓			
3	3					✓	
4	2			✓		✓	
5	2			✓	✓	✓	
6	2				✓		
7	2				✓	✓	
8	1						✓
9	1				✓		✓
10	1	✓			✓	✓	

Query result 1: Infection
 Located nearby/on/at/in → anatomical location.
 Query result 2-10: Observation
 Located nearby/on/at/in → skin.

To limit the scope of modeling the clinical knowledge ontology, 15 documents were used as examples to form the classes and individuals. Each document was split into sentences to identify catheter and infection indications at the sentence-level and document-level. At the sentence-level, individual sentences were presented to clinicians who determined what observations, anatomical locations, or procedures within the text are needed to determine catheter and infection indication. Only clinician-identified sentences with indications were included as individuals in the ontology. At the document-level, individual sentences from a document were presented together, allowing clinicians to identify indications based on additional information from a more complete documented story. Presenting the document as separate sentences allows clinicians to identify concepts within a limited example to determine what can and cannot be determined based on limited information. Whereas, allowing a clinician to see the whole document presented more possibilities and helped identify necessary data combinations for indications of catheters and infections.

Because the focus of the ontology includes catheter indications and the clinicians have identified that it is important to identify infusion phlebitis, infusion phlebitis was included in the ontology as rules based on the 1998 Visual Infusion Phlebitis Scale [56] in-

cluded in the 2021 Infusion Therapy Standards of Practice Updates [57]. Furthermore, causality is not within the scope because the exact reason for chemical and mechanical reactions resulting in infection-like signs are more likely found at the body's cellular or genetic-level in pathophysiology studies [59, 60] and unlikely to be documented in AE documents.

Anatomical locations in this ontology were kept simple and similar to the AAENOTE. Clinical guidelines for catheter insertion into anatomical locations are very specific (i.e., a central venous catheter is inserted in the jugular vein until it reaches the superior vena cava [61]) because clinical guidelines provide instructions on how to perform a task properly. However, clinical documentation is more general (i.e., central venous catheter in the chest) because this is common clinical knowledge, and the documentation is written for other clinicians to understand. To match the ontology with available documented data, this ontology relies on general anatomical location terminology. If clinicians deem it necessary, clinical guidelines can be included in a separate ontology focused on identifying catheter locations based on clinical guidelines and the FMA. Inclusion of clinical guidelines to identify specific catheter insertion sites and placement requires anatomical knowledge. For instance, to identify a central venous catheter's general anatomical location using a clinical guideline and the anatomy ontology FMA, the ontology would need to:

1. Identify the jugular vein insertion site and the superior vena cava placement.
2. Infer that the jugular vein is in the neck and the superior vena cava is present within the superior and middle mediastinum [62], annotated as anatomical location chest.
3. Convert the terms into more general terms that match the available data (i.e., vena cava is in the chest).

7.2. Representing Domain Knowledge

Discussions with clinicians about example documents and indications formulated the classes, object properties, data properties, and rules within the ontology. Then, the provided indications were sorted and summarized to match the ontology closely. Afterward, indications were verified by clinicians and included in the ontology using SPARQL queries. A list of indications can be found in Appendix C.3.1 to Appendix C.3.7.

8. Ontology Results for Domain Knowledge

8.1. Catheter Infection Indications Ontology (CIIO)

The Catheter Infection Indications Ontology (CIIO) represents clinical knowledge for signs of infections and catheters to identify PIVC-related BSIs and was developed to accompany the AAENOTE. Similar to AAENOTE, this ontology is also in both English and Norwegian. There are 57 classes, 10 object properties, 16 data properties, and 187 individuals. The 7 classes which form the main hierarchies are:

1. Observation: Any sign or symptom that can be monitored.
2. Anatomical Location: Any anatomical body part, organ, or relative position of the body.
3. Medical Device: Any instrument, device, or equipment used for a medical purpose.
4. Procedure: Any procedure, intervention, or activity related to catheters.
5. Person: An individual.
6. Document: Unstructured free-text report consisting of sentences documented to represent observable patient states.
7. Sentence: A set of words documented to represent observable patient states.

Relationships between the 7 class hierarchies can be formed using the following 10 object properties:

1. Person's subclass Patient $\xrightarrow{\text{Patient has}}$ Observation, Anatomical Location, Medical Device, or Procedure.
2. Procedure $\xrightarrow{\text{Procedure uses}}$ Medical Device.
3. Observation $\xrightarrow{\text{Is observed with}}$ Medical Device or Procedure.
4. Observation, Medical Device, or Procedure $\xrightarrow{\text{Located nearby/on/at/in}}$ Anatomical Location.
5. Document or Sentence $\xrightarrow{\text{Contains}}$ Observation, Anatomical Location, Medical Device, Procedure, or Person. Additionally, only Document $\xrightarrow{\text{Contains}}$ Sentence.
6. Procedure's subclass General IV $\xrightarrow{\text{Is combined with}}$ a different General IV
7. Procedure's subclass General IV $\xrightarrow{\text{Is IV for}}$ Procedure's subclass Infusion.

8. Procedure's subclass General IV $\xrightarrow{\text{Medication should have been}}$ a different General IV.
9. Observation, Anatomical Location, Medical Device, Procedure, or Person $\xrightarrow{\text{Is documented in}}$ Document or Sentence. Additionally, only Sentence $\xrightarrow{\text{Is documented in}}$ Document.
10. Anatomical Location $\xrightarrow{\text{Location has}}$ Observation, Medical Device, or Procedure.

An example showing how a sentence is represented using the class hierarchies and class properties of CIIO to model documented clinical knowledge is shown in Fig. 6. The complete class hierarchies of CIIO are in Appendix C.1.

Each sentence documented in the report is an individual of the Sentence class. Sentence class individuals contain Observation, Anatomical Location, Medical Device, Procedure, or Person individuals present within the text. An individual of the Document class contains the Sentence individuals that form it and the content from those sentences. Similar to AAENOTE, individuals of Observation, Anatomical Location, Medical Device, Procedure, and Person can also have object properties and data properties.

8.2. CIIO Evaluation

Designed to capture and reason about clinical catheter-related and infection-related signs and symptoms documented in an AE report, the CIIO provides the missing clinical domain knowledge for the AAENOTE. CIIO can answer 8 of the 9 competency questions based on assumptions and indications. The assumptions are that 1 AE document represents 1 patient and all sentences within a document are likely to describe concepts within the same event (Appendix C.2). Indications for catheters and infections are provided in (Appendix C.3). Additionally, the ontology classes and relationships used to answer each competency question is detailed in Appendix C.4.

9. Discussion

9.1. Ontology Development Method Comparison

The clinical problem drove this study, and the objective was not to apply an ontology development method.




Fig. 6. Catheter Infection Indications Ontology (CIIO) Clinical Knowledge Representation. (a) A sentence from a document used to identify documented clinical knowledge. Annotations are based on terms from the Annotated Adverse Event NOTE Terminology (AAENOTE). (b) AAENOTE terms were used in CIIO to conceptualize the classes and object properties that represent documented knowledge and can be used in reasoning to identify catheters and infections. Part of the ontology class hierarchies and object properties used in knowledge representation are shown. Classes are in white boxes and object properties are shown using thicker colored lines with arrow.




Fig. 7. Development phases for Annotated Adverse Event NOTE Terminology (AAENOTE) and Catheter Infection Indications Ontology (CIIO).

Hence, a specific ontology development method was not applied. However, certain steps taken are similar to the pre-existing methods and this study does include the typical phases of specification, conceptualization, formalization, implementation, maintenance, knowledge acquisition, evaluation, and documentation. An overview of the process is shown in Fig. 7, and similarities to other methods can be found in Appendix A.

During the pre-development and specification phases of the terminology, clinicians provided the research question and use case. Those were utilized to define the competency questions. Additionally, the AE dataset was retrieved, and an AE synthetic dataset was created. The conceptualization phase was performed by iteratively developing the annotation guideline and annotation sessions. Afterward, the formalization and im-

plementation phases of the terminology was developed iteratively based on the annotation guideline and using instances from the annotated corpus to answer competency questions for evaluation. Knowledge acquisition occurred during all phases with insight, guidance, and feedback from clinicians. Documentation is provided in the annotation guidelines, the annotated corpus, and the evaluation of competency questions. The annotation guidelines document changes in terms over time, and the annotated corpus documents knowledge acquisition from the text. Answers to each competency question are documented using natural language for clinicians and SPARQL queries for computer scientists.

Ontology development is similar to the terminology's pre-development, specification, and conceptualization phases. However, the formalization and implementation phases differ. The ontology iteratively incorporated clinical knowledge that can be annotated in AE documents using terminology terms to answer competency questions for evaluation. Knowledge acquisition was provided through the annotated corpus, clinician-provided catheter indication rules, and clinician-provided publications containing phlebitis rules. Additionally, clinicians iteratively reviewed and verified documented sentences to match the rules and competency questions. Ontology documentation includes the assumptions (Appendix C.2), rules for catheter and infection indications (Appendix C.3), and how competency questions were answered (Appendix C.4).

Although UMLS includes clinical terminology (i.e., SNOMED CT and ICD-10), the terms are often a combination of different concepts. For example, phlebitis has many options and is combined with different locations, such as "phlebitis of the lower limb vein," "phlebitis of the portal vein," and "retinal phlebitis." Additionally, swollen has many options, such as "foot swelling," "swollen nose," and "tongue swelling." The June 10, 2022 version of SNOMED CT has 361,907 classes⁴. It would require extensive time and effort to determine which classes are suitable for our purpose and to maintain a pre-determined class hierarchy. Introducing UMLS terminology would make it difficult for annotators to determine which term to use, introduce ambiguities for the use case, and decrease the precision needed. Furthermore, the UMLS MetaMap

is software that finds and links biomedical text to terminology concepts [23]. Unfortunately, that software is for biomedical text and not clinical text. Clinical text differs from biomedical text because it is often ungrammatical and ambiguous, with many shorthand abbreviations and acronyms [63]. Additionally, the MetaMap developers have mentioned that detecting abbreviations and acronyms needs improvements [23].

9.2. AAENOTE Scope and Limitations

The clinician asks questions about the condition of a physical patient and the patient's PIVCs, but AAENOTE is about words in the AE document regarding a patient event. The correspondence between clinical condition and document content is represented by CIO. So the answer to a question about a patient's condition will be answered from the document annotation and text. Understanding a query means translating it from clinical concepts to document content concepts. Here, the terminology is used to fit and answer questions. SPARQL queries can answer most competency questions, and the results can be used as consistency checks. For example, SPARQL can be used to count and make quantitative queries about the number of catheters and devices. Likewise, qualitative results enabled clinicians to verify if results matched their expectations of clinical events (i.e., the anatomical location of specific catheters) or why the AE was reported (i.e., incorrect medical devices used in a particular procedure).

There are several limitations to the AAENOTE. Although this terminology does not cover sepsis, it does cover events that could lead to sepsis. This terminology lacks the clinical knowledge required to answer several competency questions more in-depth. Moreover, it is not always possible to determine what a patient has because of the document's content or provided annotations. Most documents do not explicitly mention a patient because these are AE documents, and it is often implied that the adverse event has happened to a patient. Annotators will often not link the patient to all possible observations, anatomical locations, medical devices, or procedures because typically, one AE document refers to one event or patient. Furthermore, referent tracking and resolution are not handled by AAENOTE. Thus, multiple mentions of a label or individual do not indicate whether it is the same item or a different item. For example, given the annotated document in Fig. 2, the terminology cannot determine if the 2 `patient` individuals refer to the same patient

⁴<https://bioportal.bioontology.org/ontologies/SNOMEDCT/?p=summary>

1 or not because each label is an individual. Similarly,
 2 the same also applies to the 2 PIVC individuals. In this
 3 example, a query counting how many patients have a
 4 PIVC will answer 2. However, based on the context,
 5 both sentences in the document likely refer to the same
 6 patient and PIVC.

9.3. CIIO Scope and Limitations

10 The CIIO is an abstract ontology with instances
 11 populated using the terminology. Only parts of the
 12 AAENOTE necessary for creating queries with clinician-
 13 provided indications were included or extended. This
 14 provides flexibility, allows for easier ontology mainte-
 15 nance, and separates the needs of clinicians who use
 16 CIIO and annotators who use AAENOTE.

17 Based on assumptions and indications, competency
 18 questions can be answered using SPARQL queries.
 19 The queries retrieve documents and translate the con-
 20 tent for the user by identifying concepts necessary to
 21 answer the competency questions. Thus, the retrieved
 22 documents and concepts can provide sufficient infor-
 23 mation for clinicians to further decipher retrieved an-
 24 swers. For example, the exact reason why a patient
 25 needs a catheter cannot be determined by the query
 26 unless there is a direct relationship (i.e., *procedure*
 27 *Procedure uses* → *medical device*) because the list of
 28 indications does not provide reasons for catheter us-
 29 age. However, the clinician can view the retrieved list
 30 of medical devices and procedures within a document
 31 to determine why the medical devices were required.

32 The lack of detailed documentation inhibits the
 33 query's ability to answer certain questions. This in-
 34 cludes why a patient needs a catheter as previously
 35 stated, counting catheters within a patient, and where
 36 the catheters are located in a patient. Counting the ex-
 37 act number of catheters per document is not possible
 38 because multiple sentences within a document could
 39 be describing the same catheter or multiple procedures
 40 could use the same catheter. The exact anatomical lo-
 41 cation of catheters per document cannot be determined
 42 for several reasons. First, multiple sentences within a
 43 document could be describing the same catheter at the
 44 same location but with more general terms (i.e., *arm*
 45 instead of *hand*). Second, the location's position not
 46 being documented makes it difficult to distinguish if a
 47 body part is on the same side. Finally, various proce-
 48 dures can be performed at the same location. For ex-
 49 ample, given an example document, "The patient re-
 50 ceived IV fluids in elbowA and IV antibiotics in right
 51 handB. Right armC showed signs of phlebitis." Here,

1 handB is part of armC because both are on the right
 2 side, but elbowA might or might not be part of armC.
 3 Additionally, armC is likely a more general term for
 4 handB. Furthermore, an additional anatomical ontol-
 5 ogy is needed to infer the possible locations based on
 6 catheter type.

9.4. Purpose of Separate Terminology and Ontology

10 Even though the ontology uses terms from the ter-
 11 minology, the terminology and ontology are separate.
 12 They are separate because of their different purposes
 13 and functionalities. Additionally, separation provides
 14 downstream analysis flexibility for researchers. It also
 15 simplifies evaluation and allows for easier mainte-
 16 nance. Furthermore, separation enables a better under-
 17 standing of the terminology's and ontology's limita-
 18 tions.

19 The terminology and ontology were developed for
 20 different purposes using different methods. AAENOTE
 21 is intended to be useful for annotators who are annotat-
 22 ing documentation and a way to provide them a struc-
 23 tured terminology with varying granularity. In compar-
 24 ison, CIIO is intended to be used by clinicians to clin-
 25 ically reason about a patient state. The design process
 26 of AAENOTE is heavily based on the explicit terms
 27 used in annotations and not on competency questions.
 28 It uses the bottom-up method and the annotation guide-
 29 line development process to capture semantic annota-
 30 tions. In contrast, the design process of CIIO is based
 31 on the competency questions, which focus on patient
 32 states. Designed with a top-down method, it is based
 33 on concepts naturally used by clinicians to describe
 34 patients. Thus, the terminology and ontology have dif-
 35 ferent purposes and functionalities.

36 Separating the terminology from the ontology en-
 37 ables annotators to annotate concepts with standard
 38 terms and clinicians to reason about the annotated con-
 39 cepts. Here, the ontology does not impact the termi-
 40 nology annotators can use. Instead, the ontology pro-
 41 vides knowledge for the terminology. Thus, our meth-
 42 ods avoid the significant amount of time, effort, and
 43 manual curation previously required to map terms to
 44 an ontology [18]. Instead, our ontology utilizes con-
 45 cepts in the terminology and is limited by the compe-
 46 tency questions, clinical guidelines used, and clinician-
 47 provided rules. In downstream analyses, researchers
 48 can freely choose to use the terminology to quickly re-
 49 trieve documents with specific annotations, the ontol-
 50 ogy to reason and infer clinical knowledge, or both.

The terminology indicates concepts annotated in documents. Using terms from the terminology ensures that the included clinical knowledge within the ontology represents the knowledge documented in the text that can be annotated. As the ontology develops further, it is possible to conceptualize additional terms required to answer the competency questions. Those terms can then be added to the terminology and annotation guideline for additional data curation.

In this paper, separating the terminology made it easier and quicker to evaluate syntax and semantics because the ontology only has 187 instances compared to the terminology's 4470 instances. Mixing the indexed annotation terminology with a clinical knowledge ontology would be outside the ontology's scope, decrease ontology reusability, and increase the complexity of ontology maintenance. Additionally, the terminology can cover a broader scope of documents not in the ontology. Finally, using competency questions to evaluate the terminology and ontology separately reveals the distinct limitations of both. The inability to answer competency questions can be due to either the lack of knowledge or lack of necessary content within the data.

9.5. Representing Annotated Data and Revisions

Annotating data provides data meaning, and the corresponding annotation guideline and terminology provide additional structured semantic meaning. Additionally, the terminology can represent knowledge and disambiguate annotation entities and relationships. Each annotation session uses a slightly different annotation guideline that has been revised based on the previous annotation session. Hence, revisions in the annotation guidelines include added, re-organized, and removed categories. Since results from 4 different annotation sessions are included as individuals in the AAENOTE, this indicates the terminology can handle different versions of annotated data while preserving semantic meaning. It is also possible to easily customize the granularity (i.e., superclasses, classes, or subclasses) and extend or retract the terminology based on clinician needs without breaking the terminology.

To alleviate the problem with overlapping terms and ambiguities experienced by [17] and remove mismatches between the annotations and the terminology experienced by [18], all annotators in our study could only use provided annotation labels from the terminology. Using concrete concepts from the annotation guideline based on what can be found in the documen-

tion instead of other pre-existing ontologies lowers the complexity and simplifies the terminology.

9.6. Representing Annotated Documents The Way Clinicians View Patients

The AAENOTE is a terminology that provides an index of what is annotated in a clinical document. It is not used to design a language's syntax, grammar, or terms because AAENOTE is a terminology for understanding the language and underlying meanings. Instead, it is the interpreted formalized language that has been translated into basic statements for reasoning. To capture relevant information, the underlying document was represented by annotated labels and relationships for the task of question answering and text understanding instead of solely retrieving information. Hence, the terminology focuses only on items of interest and is blind to items not within the terminology.

The corresponding CIIO is an ontology that models clinical knowledge missing from AAENOTE. It provides the missing clinical knowledge required to reason about the presence of catheters and infections documented in clinical text. Although the data modeled is documented text, it enables clinicians to think about the data as an individual patient because they already do this routinely when documenting patient states.

9.7. Understandability and Accessibility for Domain Experts

The approach in this study made ontology development understandable and accessible for the domain experts without formal ontology training. Furthermore, the employed approach made it possible for clinicians to understand and be part of the design process. In practice, the approach was a necessity to progress in developing the ontology to incorporate clinical knowledge.

9.8. Clinical Utility

The collected competency questions and requirements are largely met. Thus, the main objective of developing a terminology and ontology that clinicians and hospital systems can use to get a systematic overview of identifying and reasoning about PIVC-related phlebitis and sepsis in an AE corpus has been met. Furthermore, our ontology is a step toward automated and continuous quality control that move beyond today's focus on repeated point prevalence qual-

ity controls, like the Peripheral Intravenous Catheter mini Questionnaire (*PIVC-miniQ*) [64].

The developed ontology is of value for sepsis because of its purpose, clinician involvement during development, and intended use. The ontology focuses on identifying indicators of catheter-related phlebitis or infections that can lead to sepsis by utilizing the clinicians' documentation and perspectives. Throughout the whole development, clinicians were involved as the users, domain experts, and data annotators. Furthermore, clinical knowledge within the ontology was captured similarly to how clinicians ask questions, document observations, and view documents as patients. The intent is to eventually implement the ontology into a quality surveillance system to automatically detect the presence of PIVC-related phlebitis and BSIs to improve PIVC care and lower sepsis incidents. Thus, only documented content can be included as data, and the ontology must directly correspond to and represent concepts documented within the AE documents from the clinician's perspective.

10. Future Work

For the sepsis-related use case, the synthetic AE dataset used for annotations is a placeholder for the real Norwegian AE dataset and clinical records from the EHR. Future work includes utilizing the current AAENOTE to annotate the real Norwegian AE dataset and clinical records. And to evaluate if clinical knowledge from the CIIO can still be applied and expanded on new data. Additionally, the ontologies could be applied to AE documents at other Norwegian hospitals to assess how similar documentation and knowledge are between different hospitals. The ontologies can be directly translated to other Scandinavian languages (i.e., Swedish and Danish) and applied similarly at other Scandinavian hospitals. The design and representations are largely language-independent and should be easy to transform for English clinical text about adverse events. After all, international literature suggests that the phenomena related to PIVC and devices are language-independent [3]. It would also be possible to provide multi-language querying over multi-language AE documents to enable cross-language repositories [65]. Furthermore, supervised machine learning methods can be employed to identify PIVC-related BSIs and classify patients requiring additional monitoring.

11. Conclusion

The development process resulted in a terminology and an ontology, specifically, the Annotated Adverse Event NOte Terminology (AAENOTE) which models annotated classes in annotated documents and the Catheter Infection Indications Ontology (CIIO) which models clinical knowledge for catheter and infection indications. Although there is a clinical focus here, the methodology for creating a terminology from an annotation guideline for semantically annotated data and a domain knowledge ontology to represent knowledge can be utilized in other domains to provide additional semantic meaning to annotated datasets in other domains.

12. Data Availability

The AAENOTE, CIIO, and SPARQL queries for this paper are in the GitHub repository branch "swj" at https://github.com/melissayan/aaenote_and_ciio. Detailed specifications for AAENOTE and CIIO were generated using Wizard for DOCUMENTING Ontologies (WIDOCO) [66] and are available in English and Norwegian at <https://folk.ntnu.no/melissay/ontology/index.html>.

Acknowledgements

The authors would like to acknowledge the eight annotators for producing the annotated clinical adverse event (AE) corpus. Funding was provided by the Computational Sepsis Mining and Modelling project through the Norwegian University of Science and Technology (NTNU) Health Strategic Area. Additionally, ethical approval was granted by the Regional Committees for Medical and Health Research Ethics (REK), approval no. 26814.

Appendix A. Ontology Development Methods and Evaluation Similarities

There are various ontology development methods and the steps in this study have some similarities to other methods Table 6.

1	1
2	2
3	3
4	4
5	5
6	6
7	7
8	8
9	9
10	10
11	11
12	12
13	13
14	14
15	15
16	16
17	17
18	18
19	19
20	20
21	21
22	22
23	23
24	24
25	25
26	26
27	27
28	28
29	29
30	30
31	31
32	32
33	33
34	34
35	35
36	36
37	37
38	38
39	39
40	40
41	41
42	42
43	43
44	44
45	45
46	46
47	47
48	48
49	49
50	50
51	51

Table 6
Ontology Development Methods Versus This Paper

Uschold King [31]	and	Grüniger Fox [32]	and METHONTOLOGY [33]	On-To-Knowledge [34]	NeOn [35]	This Paper
Pre-development				Feasibility study phase to identify problems.	Initiation phase to specify ontology requirements, intended use, users, and formal language in the form of competency questions and glossary of terms.	Use clinician identified research question to form competency questions. Form terms for annotation, perform data screening and pre-annotation, develop annotation guideline, schedule annotation sessions, and evaluate annotations.
Specification		Define ontology requirements as competency questions based on a scenario or user provided problem.	Specification phase to produce ontology specifications.	Kickoff phase to document ontology requirements and semi-formal ontology.	Part of pre-development.	Part of pre-development.
Conceptualization		Define terminology of ontology.	Conceptualization phase to build glossary of terms.	Refinement phase to refine semi-formal ontology and formalize ontology iteratively based on domain expert interviews.	Design phase to produce an informal and formal model to meet requirements.	Annotation and annotation guideline revision phases determine the terms and concepts that are documented in data.
Formalization		Specify terminology definitions using first order logic.	Part of conceptualization.	Part of conceptualization.	Implementation phase to implement the formal model into an ontology language.	Codify terminology and ontology in Protégé using OWL.
Implementation		Represent terminology in a formal language.	Implementation phase to codify ontology into formal ontology.			Part of formalization.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

Table 6
(continued)

	Uschold and King [31]	and Grüninger and Fox [32]	METHONTOLOGY [33]	On-To-Knowledge [34]	NeOn [35]	This Paper
Management			Interview experts and/or analyze text.	Apply the ontology and manage its evolution and maintenance.	Use the ontology to detect errors or missing knowledge for design phase of the next ontology version.	Terminology can be updated to a new annotation guideline version and its corresponding annotated corpus. Ontology can include new clinical knowledge.
Knowledge Acquisition				Knowledge creation, capture, retrieval and access, and use.	Knowledge is introduced by domain experts and ontology practitioners at different development phases.	Knowledge was acquired iteratively with users and domain experts. Discussions with users and domain experts resulted in annotation guideline revisions and discovering the knowledge needed for ontology reasoning and inferences. Annotations by domain expert annotators to capture knowledge from text.
Evaluation	Adapt what has been done in the knowledge based systems field for ontologies.	Evaluate the ontology by proving completeness theorems to answer competency questions.	Evaluate the ontologies, software environment, and documentation with the requirement specification document during and between each phase. Document how the ontology was evaluated, errors detected, and knowledge sources for evaluation.	Proposed technology-focused (i.e., the development tool's evaluation of syntax and semantics of the ontology and the evaluation of tools and applications for interoperability and scalability), user-focused (i.e., user satisfaction with the application and comparing the ontology based application to pre-existing ones), and formal evaluation.	Evaluate the ontology using 5 different tasks [67] which includes selecting individual ontology network components for evaluation, selecting the evaluation goal and approach, identifying the reference and evaluation metric, applying the selected evaluation, and presenting the combined results of the individual components.	Ability to answer competency questions and if the ontology is useful from the perspective of the user.
Documentation	Document ontology type and purpose		Document ideally all phases, knowledge acquisition, and evaluation.	Document kickoff phase ontology requirements.	Document ontology requirements, ontology description, and evaluation.	Annotation guidelines record changes, annotations document knowledge acquisition, and evaluation documentation answers competency questions in a format understandable to the users and developers.

Appendix B. Annotated Adverse Event Note Terminology (AAENOTE) Hierarchy Competency Questions (CQs), SPARQL Queries, and Results

B.1. AAENOTE Hierarchy

The AAENOTE classes and their subclasses can be found in the following figures:

1. Fig. 8 Observation
2. Fig. 9 Anatomical Location
3. Fig. 10 Medical Device
4. Fig. 11 Procedure
5. Fig. 12 Identifier
6. Fig. 13 Person
7. Fig. 14 Annotated Document




Fig. 8. AAENOTE Observation Class Hierarchy




Fig. 9. AAENOTE Anatomical Location Class Hierarchy




Fig. 10. AAENOTE Medical Device Class Hierarchy




Fig. 11. AAENOTE Procedure Class Hierarchy




Fig. 12. AAENOTE Identifier Class Hierarchy




Fig. 13. AAENOTE Person Class Hierarchy




Fig. 14. AAENOTE Annotated Document Class Hierarchy

B.2. Details About Converting Annotation Guideline to Terminology

To further differentiate observations, Descriptive sign or symptom, Vital sign, Neurological and physiological, Wound, Lab result, and Device observation subclasses were introduced. In addition, Insertion site was introduced because some documents document a catheter’s insertion site without mentioning a specific body part. Similarly, Organ parts, such as Skin and Circulatory system, were also introduced because they are documented instead of body part. For Medical Device, additional catheters were included (i.e., Intraosseous cannula and subcutaneous catheter for specificity. The category for device parts was removed because the terminology only covers catheter and catheter parts; thus, it is a part of a medical device if it is not under a catheter. For Procedure, Administration way was introduced to describe how a substance was administered into the patient, Catheter procedure subclasses all include “catheter” to indicate the action is for catheters only, and different Infusion types were included to differentiate from Injection types.

A hierarchy for certain data properties was introduced to organize the anatomical location descriptors, contents within a document, and document identity. For instance, document identity-related data properties were included for identification, such as annotator ID, filename, and annotation session. Unlike the data property values for body temperature (i.e., hyperthermia, normal, and hypothermia) and severity level (i.e., high, normal, and low), Consciousness level required a separate data property with values of alert, confusion, painfully responsive, unresponsive, and verbally responsive). The “observation is diagnosed by” and “observation said by” data properties were added to distinguish signs from symptoms because signs are what a clinician observes and symptoms are what a patient says.

B.3. AAENOTE Competency Questions and Terminology Usage

Table 7: AAENOTE Competency Questions and Terminology Usage

Competency Question	Terminology Classes and Relationships to Find Instances*
1. Does patientA have phlebitis, and was it infectious phlebitis, chemical phlebitis, or mechanical phlebitis?	<p>Explicit:</p> <p>Patient $\xrightarrow{\text{Person has}}$ phlebitis.</p> <p>Implicit:</p> <p>Phlebitis $\xrightarrow{\text{Located nearby/on/at/in}}$ anatomical location. [†]Anatomical location is part of a person, and typically a patient given the context is an AE note. [‡]Need clinical knowledge to determine if the phlebitis is infectious phlebitis, chemical phlebitis, or mechanical phlebitis.</p>

1	2. Does patientA have an infection?	Explicit:	1
2		Patient $\xrightarrow{\text{Person has}}$ infection.	2
3		Patient $\xrightarrow{\text{Person has}}$ infection and other observation (s).	3
4		Implicit:	4
5		Infection $\xrightarrow{\text{Located nearby/on/at/in}}$ anatomical location.	5
6		Infection and other observation (s) $\xrightarrow{\text{Located nearby/on/at/in}}$	6
7		anatomical location.	7
8		†Anatomical location is part of a person, and typically a pa-	8
9		tient given the context is an AE note.	9
10		‡Need clinical knowledge to determine if the observations	10
11		combined indicate infection without infection explicitly in-	11
12		cluded.	12
13	3. Does patientA have a BSI?	§ Cannot determine if there is a BSI without a microbiology laboratory	13
14		result of a positive blood culture and the cultured bacteria name.	14
15	4. How many patients have an infection or BSI?	Same as Competency Question 1 and 2.	15
16			16
17	5. Which patients have sepsis?	Explicit:	17
18		Patient $\xrightarrow{\text{Person has}}$ sepsis.	18
19		Patient $\xrightarrow{\text{Person has}}$ sepsis and other observation (s).	19
20		Implicit:	20
21		‡Need clinical knowledge to determine if the observations	21
22		combined indicate sepsis.	22
23			23
24	6. Does patientB have a catheter?	Explicit:	24
25		Patient $\xrightarrow{\text{Person has}}$ catheter.	25
26		Implicit:	26
27		Catheter $\xrightarrow{\text{Located nearby/on/at/in}}$ anatomical location.	27
28		†Anatomical location is part of a person, and typically a pa-	28
29		tient given the context is an AE note.	29
30		‡Need clinical knowledge to determine if the observations,	30
31		anatomical locations, and/or procedures combined in-	31
32		indicate a catheter is present.	32
33			33
34	7. Does patientB have a PIVC?	Explicit:	34
35		Patient $\xrightarrow{\text{Person has}}$ PIVC.	35
36		Implicit:	36
37		PIVC $\xrightarrow{\text{Located nearby/on/at/in}}$ anatomical location.	37
38		†Anatomical location is part of a person, and typically a pa-	38
39		tient given the context is an AE note.	39
40		‡Need clinical knowledge to determine if the observations,	40
41		anatomical locations, and/or procedures combined in-	41
42		indicate a PIVC is present.	42
43			43
44	8a. How many catheters does patientB have?	Same as Competency Question 5.	44
45			45
46			46
47			47
48			48
49			49
50			50
51			51

1 8b. Where are the catheters in patientB?

Explicit:

Patient $\xrightarrow{\text{Person has}}$ anatomical location (X).

Patient $\xrightarrow{\text{Person has}}$ Catheter (Y).

Catheter (Y) $\xrightarrow{\text{Located nearby/on/at/in}}$ anatomical location (X).

Implicit:

Catheter $\xrightarrow{\text{Located nearby/on/at/in}}$ anatomical location.

†Anatomical location is part of a person, and typically a patient given the context is an AE note.

11 8c. Why does patientB need the catheter(s)?

Explicit:

(a) Patient has a catheter and the catheter is used in the procedure.

Patient (X) $\xrightarrow{\text{Person has}}$ catheter (Y).

Procedure $\xrightarrow{\text{Procedure uses}}$ catheter (Y).

(b) Patient has a catheter, patient has a procedure, and that procedure uses that catheter.

Patient (X) $\xrightarrow{\text{Person has}}$ catheter (Y).

Patient (X) $\xrightarrow{\text{Person has}}$ procedure (Z).

Procedure (Z) $\xrightarrow{\text{Procedure uses}}$ catheter (Y).

Implicit:

(a) Patient has a catheter and patient has a procedure.

Patient (X) $\xrightarrow{\text{Person has}}$ catheter.

Patient (X) $\xrightarrow{\text{Person has}}$ procedure.

‡Need clinical knowledge to determine which procedure is likely to use or involve a specific type of catheter.

29 9a. Does patientC have an infection and catheter?

Explicit:

Patient (X) $\xrightarrow{\text{Person has}}$ infection.

Patient (X) $\xrightarrow{\text{Person has}}$ catheter.

Implicit:

‡Need clinical knowledge to determine if the observations combined indicate infection.

‡Need clinical knowledge to determine if the observations, anatomical locations, and/or procedures combined indicate a catheter is present.

39 9b. Was patientC's infection associated with a catheter?

§ Cannot determine if an infection is associated with a catheter unless that catheter is tested in the microbiology lab.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

B.4. AAENOTE CQ 1: Does patientA have phlebitis, and was it infectious phlebitis, chemical phlebitis, or mechanical phlebitis?

The patients who have phlebitis are listed in Table 8 using Listing 1. And anatomical locations with phlebitis in Table 9 were queried using Listing 2.

```

PREFIX rdf: <http://www.w3.org/1999/02/22-rdf-syntax-ns#>
PREFIX owl: <http://www.w3.org/2002/07/owl#>
PREFIX rdfs: <http://www.w3.org/2000/01/rdf-schema#>
PREFIX xsd: <http://www.w3.org/2001/XMLSchema#>
PREFIX : <http://www.semanticweb.org/2022/04/aaenote#>
SELECT DISTINCT ?indv_patient ?class_phlebitis
WHERE {
  ?indv_patient rdf:type :patient ;
               rdf:type ?class_patient .
  FILTER(?class_patient != owl:NamedIndividual) .
  ?indv_phlebitis rdf:type* :phlebitis ;
                 rdf:type ?class_phlebitis .
  FILTER(?class_phlebitis != owl:NamedIndividual) .
  FILTER(REGEX(str(?class_phlebitis), "phlebitis")) .
  ?indv_patient :person_has ?indv_phlebitis .
}
ORDER BY ?indv_patient
    
```

Listing 1: Patient has phlebitis.

Table 8
AAENOTE SPARQL Query Result: List patients that have phlebitis

Query Result	Patient	Phlebitis
1	patient.T1.2.lo22a.SD_0003	✓
2	patient.T1.3.do13a.PO_0005	✓
3	patient.T2.2.do13a.SP_0008	✓
4	patient.T2.2.lo12a.SD_0003	✓
5	patient.T2.2.lo22a.PO_0005	✓
6	patient.T2.3.do23a.PO_0005	✓
7	patient.T2.3.po24a.SD_0003	✓
8	patient.T2.4.lo22a.SP_0008	✓
9	patient.T3.3.po14a.SD_0003	✓
10	patient.T4.1.po14a.PO_0005	✓

Query result 1-10: Patient $\xrightarrow{\text{Person has}}$ phlebitis.

```

1 PREFIX rdf: <http://www.w3.org/1999/02/22-rdf-syntax-ns#>
2 PREFIX owl: <http://www.w3.org/2002/07/owl#>
3 PREFIX rdfs: <http://www.w3.org/2000/01/rdf-schema#>
4 PREFIX xsd: <http://www.w3.org/2001/XMLSchema#>
5 PREFIX : <http://www.semanticweb.org/2022/04/aaenote#>
6 SELECT DISTINCT ?indv_location ?class_phlebitis
7 WHERE {
8   ?indv_phlebitis rdf:type :phlebitis ;
9     rdf:type ?class_phlebitis .
10  FILTER(?class_phlebitis != owl:NamedIndividual) .
11  ?indv_location rdf:type/rdfs:subClassOf* :anatomical_location ;
12     rdf:type ?class_location .
13  FILTER(?class_location != owl:NamedIndividual) .
14  ?indv_phlebitis :located_nearby_on_at_in ?indv_location .
15 }
16 ORDER BY ?indv_location

```

Listing 2: List phlebitis located nearby/on/at/in an anatomical location.

Table 9
AAENOTE SPARQL Query Result: List phlebitis at an anatomical location

Query Result	Anatomical Location	Phlebitis
1	elbow.T1.4.lo22a.DO_0010	✓
2	elbow.T2.1.do13a.DO_0010	✓
3	elbow.T2.4.lo12a.DO_0010	✓
4	elbow.T3.3.so11a.DO_0010	✓
5	elbow.T3.3.so21a.DO_0010	✓
6	hand.T3.4.lo12a.SP_0008	✓
7	hand.T4.3.do13a.PO_0005	✓
8	hand.T7.2.lo22a.PO_0005	✓
9	wrist.T4.2.do13a.SP_0008	✓
10	patient.T4.1.po14a.PO_0005	✓

Query result 1-10: Listing 5 SPARQL query: Phlebitis
Located nearby/on/at/in anatomical location.

B.5. AAENOTE CQ 2: Does patientA have an infection?

B.5.1. AAENOTE CQ 2 Explicit

The patients who have infection are listed in Table 10 using Listing 3. Whereas, patients who have infection and/or other observations are listed in Table 11 using Listing 4.

```

PREFIX rdf: <http://www.w3.org/1999/02/22-rdf-syntax-ns#>
PREFIX owl: <http://www.w3.org/2002/07/owl#>
PREFIX rdfs: <http://www.w3.org/2000/01/rdf-schema#>
PREFIX xsd: <http://www.w3.org/2001/XMLSchema#>
PREFIX : <http://www.semanticweb.org/2022/04/aaenote#>
SELECT DISTINCT ?indv_patient ?class_infection
WHERE {
  ?indv_patient rdf:type :patient ;
               rdf:type ?class_patient .
  FILTER(?class_patient != owl:NamedIndividual) .
  ?indv_infection rdf:type+ :infection ;
                 rdf:type ?class_infection .
  FILTER(?class_infection != owl:NamedIndividual) .
  FILTER(Regex(str(?class_infection), "infection")) .
  ?indv_patient :person_has ?indv_infection .
}
ORDER BY ?indv_patient
    
```

Listing 3: List patients who have infection.

Table 10
AAENOTE SPARQL Query Result: List patients that have infection

Query Result	Patient	Infection
1	patient.T1.2.lo22a.SD_0006	✓
2	patient.T1.4.lo22a.SP_0002	✓
3	patient.T10.2.do23a.SP_0007	✓
4	patient.T12.4.lo12a.SP_0007	✓
5	patient.T2.2.do23a.SP_0006	✓
6	patient.T2.2.lo12a.SD_0006	✓
7	patient.T2.3.do23a.PO_0010	✓
8	patient.T2.3.po14a.SD_0006	✓
9	patient.T3.2.do13a.SP_0007	✓
10	patient.T4.4.lo22a.DO_0008	✓
11	patient.T4.4.so11a.DP_0010	✓
12	patient.T5.2.po24a.DO_0008	✓
13	patient.T5.4.lo22a.SP_0007	✓
14	patient.T7.3.po24a.SD_0006	✓

Query result 1-14: Patient $\xrightarrow{\text{Person has}}$ infection.

```

1 PREFIX rdf: <http://www.w3.org/1999/02/22-rdf-syntax-ns#>
2 PREFIX owl: <http://www.w3.org/2002/07/owl#>
3 PREFIX rdfs: <http://www.w3.org/2000/01/rdf-schema#>
4 PREFIX xsd: <http://www.w3.org/2001/XMLSchema#>
5 PREFIX : <http://www.semanticweb.org/2022/04/aaenote#>
6 SELECT DISTINCT ?indv_patient ?observations
7 WHERE {
8     {
9         # Group ?class_observations alphabetically by ?indv_patient
10        SELECT ?indv_patient ?class_patient
11        (GROUP_CONCAT (DISTINCT str(?class_observation); separator=", " ) as ?observations)
12        WHERE {
13            # Organize alphabetically ?class_observations
14            SELECT ?indv_patient ?class_patient ?class_observation
15            WHERE {
16                ?indv_patient rdf:type :patient ;
17                rdf:type ?class_patient .
18                FILTER(?class_patient != owl:NamedIndividual) .
19                FILTER(REGEX(str(?class_patient), "patient")) .
20                ?indv_observation rdf:type/rdfs:subClassOf* :observation ;
21                rdf:type ?class_observation .
22                FILTER(?class_observation != owl:NamedIndividual) .
23                ?indv_patient :person_has ?indv_observation .
24            }
25            GROUP BY ?indv_patient ?class_patient ?class_observation
26            ORDER BY ?class_observation
27        }
28        GROUP BY ?indv_patient ?class_patient
29    }
30    FILTER REGEX(str(?observations), "infection")
31 }
32 ORDER BY ?indv_patient

```

Listing 4: List patients who have infection and/or other observations.

Table 11
AAENOTE SPARQL Query Result: List patients that have infection and/or another observation

Query Result	Person	Observation											
	Patient	blood pressure	body temp	c reactive protein	consciousness level	infection	neurological and physiological	observation	pulse	red	respiratory rate	sepsis	swollen
1	patient.T1.2.lo22a.SD_0006	✓	✓	✓		✓			✓				
2	patient.T1.4.lo22a.SP_0002		✓			✓					✓		
3	patient.T10.2.do23a.SP_0007				✓	✓							
4	patient.T12.4.lo12a.SP_0007				✓	✓		✓					
5	patient.T2.2.do23a.SP_0006					✓							
6	patient.T2.2.lo12a.SD_0006		✓	✓		✓							
7	patient.T2.3.do23a.PO_0010					✓				✓			
8	patient.T2.3.po14a.SD_0006	✓	✓	✓		✓	✓	✓	✓		✓		
9	patient.T3.2.do13a.SP_0007			✓	✓	✓		✓					
10	patient.T4.4.lo22a.DO_0008					✓				✓			✓
11	patient.T4.4.so11a.DP_0010					✓					✓		
12	patient.T5.2.po24a.DO_0008					✓					✓		✓
13	patient.T5.4.lo22a.SP_0007			✓		✓	✓						
14	patient.T7.3.po24a.SD_0006	✓	✓	✓		✓			✓			✓	

Query result 1-14: Patient $\xrightarrow{\text{Person has}}$ infection and other observation (s).

B.5.2. AAENOTE CQ 2 Implicit

By using anatomical locations it is possible to implicitly identify infection within a patient because the anatomical locations refer to a place on a human and in the context of AE notes anatomical locations commonly refer to a place on a patient. As shown in Table 12, only 1 instance where infection is located at an anatomical location was found using either Listing 5 or Listing 6.

```

PREFIX rdf: <http://www.w3.org/1999/02/22-rdf-syntax-ns#>
PREFIX owl: <http://www.w3.org/2002/07/owl#>
PREFIX rdfs: <http://www.w3.org/2000/01/rdf-schema#>
PREFIX xsd: <http://www.w3.org/2001/XMLSchema#>
PREFIX : <http://www.semanticweb.org/2022/04/aaenote#>
SELECT DISTINCT ?indv_location ?class_infection
WHERE {
  ?indv_infection rdf:type :infection ;
                 rdf:type ?class_infection .
  FILTER(?class_infection != owl:NamedIndividual) .
  ?indv_location rdf:type/rdfs:subClassOf* :anatomical_location ;
                 rdf:type ?class_location .
  FILTER(?class_location != owl:NamedIndividual) .
  ?indv_infection :located_nearby_on_at_in ?indv_location .
}
ORDER BY ?indv_location

```

Listing 5: List infection located nearby/on/at/in an anatomical location.

```

PREFIX rdf: <http://www.w3.org/1999/02/22-rdf-syntax-ns#>
PREFIX owl: <http://www.w3.org/2002/07/owl#>
PREFIX rdfs: <http://www.w3.org/2000/01/rdf-schema#>
PREFIX xsd: <http://www.w3.org/2001/XMLSchema#>
PREFIX : <http://www.semanticweb.org/2022/04/aaenote#>
SELECT DISTINCT ?indv_location ?observations
WHERE {
  {
    SELECT ?indv_location ?class_location
      (GROUP_CONCAT (DISTINCT ?class_observation; separator=", " ) as ?observations)
    WHERE {
      {
        SELECT ?indv_location ?class_location ?class_observation
        WHERE {
          ?indv_observation rdf:type/rdfs:subClassOf* :observation ;
                           rdf:type ?class_observation .
          FILTER(?class_observation != owl:NamedIndividual) .
          ?indv_location rdf:type/rdfs:subClassOf* :anatomical_location ;
                           rdf:type ?class_location .
          FILTER(?class_location != owl:NamedIndividual) .
          ?indv_observation :located_nearby_on_at_in ?indv_location .
        }
        GROUP BY ?indv_location ?class_location ?class_observation
        ORDER BY ?class_observation
      }
    }
    GROUP BY ?indv_location ?class_location
    ORDER BY ?class_location
  }
  FILTER REGEX(str(?observations), "infection")
}
ORDER BY ?indv_location

```

Listing 6: List infection and other observations located nearby/on/at/in an anatomical location.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

Table 12
AAENOTE SPARQL Query Result: List infections and/or observations at an anatomical location

Query Result	Anatomical Location	Infection
1	skin.T4.2.po14a.SL_0006	✓

Same results for Listing 5 and Listing 6 SPARQL query.
 Query result 1 from Listing 5 SPARQL query:
 Infection Located nearby/on/at/in anatomical location.
 Query result 1 from Listing 6 SPARQL query:
 Infection and other observation(s) Located nearby/on/at/in anatomical location.

B.6. AAENOTE CQ 3: Does patientA have a BSI?

Cannot determine if there is a BSI without a microbiology laboratory result of a positive blood culture and the cultured bacteria name.

B.7. AAENOTE CQ 4: How many patients have an infection or BSI?

Explicitly, Table 4 shows the number of patients with infection and the number of patients with infection and/or other observations (Listing 7). Implicitly, there was only 1 anatomical location that had infection (Appendix B.5.2), so the results are not shown for the query in Listing 8. If there was clinical knowledge, it could provide the insight required to determine if combinations of observations at certain anatomical locations are indications of an infection (Listing 9 and Table 13). As previously stated in Appendix B.6, BSI cannot be determined because this requires microbiology laboratory blood test results.

```

PREFIX rdf: <http://www.w3.org/1999/02/22-rdf-syntax-ns#>
PREFIX owl: <http://www.w3.org/2002/07/owl#>
PREFIX rdfs: <http://www.w3.org/2000/01/rdf-schema#>
PREFIX xsd: <http://www.w3.org/2001/XMLSchema#>
PREFIX : <http://www.semanticweb.org/2022/04/aaenote#>
SELECT ?class_patient ?observations ?n_patients
WHERE {
  {
    SELECT ?class_patient
      (GROUP_CONCAT (DISTINCT str(?class_observation); separator=", " ) as ?observations)
      (COUNT (DISTINCT ?indv_patient ) as ?n_patients)
    WHERE {
      ?indv_patient rdf:type :patient ;
        rdf:type ?class_patient .
      FILTER(?class_patient != owl:NamedIndividual) .
      ?indv_infection rdf:type :infection ;
        rdf:type ?class_observation .
      FILTER(?class_observation != owl:NamedIndividual) .
      ?indv_patient :person_has ?indv_infection .
    }
    GROUP BY ?class_patient ?class_observation
    ORDER BY ?class_observation
  }
  UNION
  {
    SELECT ?class_patient ?observations (COUNT (DISTINCT ?indv_patient ) as ?n_patients)
    WHERE {
      {
        SELECT ?indv_patient ?class_patient
          (GROUP_CONCAT (
            DISTINCT str(?class_observation); separator=", " ) as ?observations)
        WHERE {
          SELECT ?indv_patient ?class_patient ?class_observation
          WHERE {
            ?indv_patient rdf:type :patient ;
              rdf:type ?class_patient .
            FILTER(?class_patient != owl:NamedIndividual) .
            ?indv_observation rdf:type/rdfs:subClassOf* :observation ;
              rdf:type ?class_observation .
            FILTER(?class_observation != owl:NamedIndividual) .
            ?indv_patient :person_has ?indv_observation .
          }
          GROUP BY ?indv_patient ?class_patient ?class_observation
          ORDER BY ?class_observation
        }
        GROUP BY ?indv_patient ?class_patient
      }
      FILTER REGEX(str(?observations), "infection")
    }
    GROUP BY ?class_patient ?observations
  }
}

```

Listing 7: Count the number of patients with infection and number patients with infection and other observations.

```

1 PREFIX rdf: <http://www.w3.org/1999/02/22-rdf-syntax-ns#>
2 PREFIX owl: <http://www.w3.org/2002/07/owl#>
3 PREFIX rdfs: <http://www.w3.org/2000/01/rdf-schema#>
4 PREFIX xsd: <http://www.w3.org/2001/XMLSchema#>
5 PREFIX : <http://www.semanticweb.org/2022/04/aaenote#>
6 SELECT ?class_location ?observations (COUNT (DISTINCT ?indv_location) as ?n_instances)
7 WHERE {
8   {
9     SELECT ?indv_location ?class_location
10    (GROUP_CONCAT (DISTINCT ?class_observation; separator=", " ) as ?observations)
11    WHERE {
12      SELECT ?indv_location ?class_location ?class_observation
13      WHERE {
14        ?indv_observation rdf:type/rdfs:subClassOf* :observation ;
15        rdf:type ?class_observation .
16        FILTER(?class_observation != owl:NamedIndividual) .
17        ?indv_location rdf:type/rdfs:subClassOf* :anatomical_location ;
18        rdf:type ?class_location .
19        FILTER(?class_location != owl:NamedIndividual) .
20        ?indv_observation :located_nearby_on_at_in ?indv_location .
21      }
22      GROUP BY ?indv_location ?class_location ?class_observation
23      ORDER BY ?class_observation
24    }
25    GROUP BY ?indv_location ?class_location
26    ORDER BY ?class_location
27  }
28  FILTER REGEX(str(?observations), "infection")
29 }
30 GROUP BY ?class_location ?observations
31 ORDER BY ?class_location

```

Listing 8: Count the number of anatomical locations where infection and other observations are located nearby/on/at/in that anatomical location.

```

1 PREFIX rdf: <http://www.w3.org/1999/02/22-rdf-syntax-ns#>
2 PREFIX owl: <http://www.w3.org/2002/07/owl#>
3 PREFIX rdfs: <http://www.w3.org/2000/01/rdf-schema#>
4 PREFIX xsd: <http://www.w3.org/2001/XMLSchema#>
5 PREFIX : <http://www.semanticweb.org/2022/04/aaenote#>
6 SELECT ?class_location ?observations (COUNT (DISTINCT ?indv_location) as ?n_instances)
7 WHERE {
8   {
9     SELECT ?indv_location ?class_location
10    (GROUP_CONCAT (DISTINCT ?class_observation; separator=", " ) as ?observations)
11    WHERE {
12      SELECT ?indv_location ?class_location ?class_observation
13      WHERE {
14        ?indv_observation rdf:type/rdfs:subClassOf* :observation ;
15        rdf:type ?class_observation .
16        FILTER(?class_observation != owl:NamedIndividual) .
17        ?indv_location rdf:type/rdfs:subClassOf* :anatomical_location ;
18        rdf:type ?class_location .
19        FILTER(?class_location != owl:NamedIndividual) .
20        ?indv_observation :located_nearby_on_at_in ?indv_location .
21      }
22      GROUP BY ?indv_location ?class_location ?class_observation
23      ORDER BY ?class_observation
24    }
25    GROUP BY ?indv_location ?class_location
26    ORDER BY ?class_location
27  }
28 }
29 GROUP BY ?class_location ?observations
30 ORDER BY ?class_location

```

Listing 9: Count the number of anatomical locations with the same observations located nearby/on/at/in that anatomical location.

Table 13

AAENOTE SPARQL Results for Anatomical Location with Combinations of Observations

Query Result	Instances	Anatomical Location				Observation										
		arm	elbow	hand	skin	body temp	edema	infection	mobility impairment	pain	phlebitis	purulence	red	swollen	warm	wound
1	12	✓								✓						
2	7	✓												✓		
3	6	✓										✓		✓		
4	4	✓								✓				✓		
5	3	✓								✓			✓	✓		
6	2	✓								✓			✓			
7	2	✓											✓			
8	1	✓								✓	✓					
9	1	✓								✓	✓		✓			
10	1	✓				✓			✓	✓		✓				
11	1	✓										✓				
12	1	✓										✓			✓	
13	1	✓					✓					✓				
14	1	✓										✓	✓	✓		
15	7		✓												✓	
16	7		✓									✓				
17	7		✓									✓	✓	✓		
18	6		✓							✓		✓	✓			
19	5		✓							✓			✓			
20	4		✓													
21	3		✓											✓		
22	2		✓									✓			✓	
23	1		✓							✓						
24	1		✓									✓				
25	4			✓										✓		
26	3			✓							✓					
27	2			✓								✓	✓		✓	
28	1			✓						✓						
29	1			✓						✓				✓		
30	1			✓								✓	✓			
31	3				✓					✓						
32	3				✓									✓		
33	2				✓					✓				✓		
34	2				✓					✓			✓	✓		
35	2				✓								✓			
36	2				✓								✓	✓		
37	1				✓										✓	
38	1				✓						✓					
39	1				✓							✓		✓		
40	1				✓					✓			✓			

Query result 1-40: Observation (s) Located nearby/on/at/in anatomical location.

B.8. AAENOTE CQ 5: Which patients have sepsis?

Explicitly, 5 patients have sepsis (Table 14, Listing 10) and 3 of the 5 patients with sepsis have sepsis and another observation (Table 15, Listing 11). Additional clinical knowledge is needed to determine if other observations combined without the sepsis class are indications of sepsis.

```

PREFIX rdf: <http://www.w3.org/1999/02/22-rdf-syntax-ns#>
PREFIX owl: <http://www.w3.org/2002/07/owl#>
PREFIX rdfs: <http://www.w3.org/2000/01/rdf-schema#>
PREFIX xsd: <http://www.w3.org/2001/XMLSchema#>
PREFIX : <http://www.semanticweb.org/2022/04/aaenote#>
SELECT ?indv_patient ?class_sepsis
WHERE {
  ?indv_patient rdf:type :patient .
  ?indv_sepsis rdf:type :sepsis ;
              rdf:type ?class_sepsis .
  FILTER(?class_sepsis != owl:NamedIndividual) .
  ?indv_patient :person_has ?indv_sepsis .
}
ORDER BY ?indv_patient
    
```

Listing 10: List patients who have sepsis.

Table 14
AAENOTE SPARQL Query Result: List patients that have sepsis

Query Result	Patient	Sepsis
1	patient.T1.2.do13a.SP_0006	✓
2	patient.T1.4.lo12a.SP_0006	✓
3	patient.T6.3.lo22a.DP_0010	✓
4	patient.T7.2.po24a.SL_0004	✓
5	patient.T7.3.po24a.SD_0006	✓

Query result 1-5: Patient $\xrightarrow{\text{Person has}}$ sepsis.

Table 15
AAENOTE SPARQL Query Result: List patients that have sepsis and/or another observation

Query Result	Person	Observation								
	Patient	blood pressure	body temp	c reactive protein	infection	pulse	red	respiratory rate	sepsis	swollen
1	patient.T1.2.do13a.SP_0006								✓	
2	patient.T1.4.lo12a.SP_0006								✓	
3	patient.T6.3.lo22a.DP_0010						✓	✓		
4	patient.T7.2.po24a.SL_0004		✓				✓		✓	✓
5	patient.T7.3.po24a.SD_0006	✓	✓	✓	✓	✓			✓	

Query result 1-5: Patient $\xrightarrow{\text{Person has}}$ sepsis and other observation (s).

```

16 PREFIX rdf: <http://www.w3.org/1999/02/22-rdf-syntax-ns#>
17 PREFIX owl: <http://www.w3.org/2002/07/owl#>
18 PREFIX rdfs: <http://www.w3.org/2000/01/rdf-schema#>
19 PREFIX xsd: <http://www.w3.org/2001/XMLSchema#>
20 PREFIX : <http://www.semanticweb.org/2022/04/aaenote#>
21 SELECT ?indv_patient ?observations
22 WHERE {
23   {
24     SELECT ?indv_patient ?class_patient
25     (GROUP_CONCAT (DISTINCT str(?class_observation); separator=", " ) as ?observations)
26     WHERE {
27       {
28         SELECT ?indv_patient ?class_patient ?class_observation
29         WHERE {
30           ?indv_patient rdf:type :patient ;
31           rdf:type ?class_patient .
32           FILTER(?class_patient != owl:NamedIndividual) .
33           ?indv_observation rdf:type/rdfs:subClassOf* :observation ;
34           rdf:type ?class_observation .
35           FILTER(?class_observation != owl:NamedIndividual) .
36           ?indv_patient :person_has ?indv_observation .
37         }
38         GROUP BY ?indv_patient ?class_patient ?class_observation
39         ORDER BY ?class_observation
40       }
41     }
42     GROUP BY ?indv_patient ?class_patient
43   }
44   FILTER REGEX(str(?observations), "sepsis")
45 }
46 ORDER BY ?indv_patient

```

Listing 11: List patients that have sepsis and/or other observations.

B.9. AAENOTE CQ 6: Does patientB have a catheter?

The patient and the type of catheter a patient has can be found explicitly using Listing 12. A subset of the results are in Table 16, where each patient individual is listed with a type of catheter and how many catheters of that specific type are present. Thus, the same patient can be listed multiple times as seen in Table 16's Query Result 28-30 where the same patient is listed 3 times because the patient has 3 catheters of different types. Implicitly, an anatomical location with a catheter indicates a person has a catheter (Listing 13, Table 17). Additional clinical knowledge is needed to determine if other observations combined indicate a catheter is present.

```

PREFIX rdf: <http://www.w3.org/1999/02/22-rdf-syntax-ns#>
PREFIX owl: <http://www.w3.org/2002/07/owl#>
PREFIX rdfs: <http://www.w3.org/2000/01/rdf-schema#>
PREFIX xsd: <http://www.w3.org/2001/XMLSchema#>
PREFIX : <http://www.semanticweb.org/2022/04/aaenote#>
SELECT ?indv_patient ?class_catheter (COUNT (DISTINCT ?indv_catheter ) as ?n_catheters)
WHERE {
  ?indv_patient rdf:type :patient ;
               rdf:type ?class_patient .
  FILTER(?class_patient != owl:NamedIndividual) .
  ?indv_catheter rdf:type/rdfs:subClassOf* :catheter ;
                rdf:type ?class_catheter .
  FILTER(?class_catheter != owl:NamedIndividual) .
  ?indv_patient :person_has ?indv_catheter .
}
GROUP BY ?indv_patient ?class_catheter
ORDER BY ?indv_patient ?class_catheter

```

Listing 12: List patients that have a catheter, the catheter's type, and the number of that catheter type.

```

PREFIX rdf: <http://www.w3.org/1999/02/22-rdf-syntax-ns#>
PREFIX owl: <http://www.w3.org/2002/07/owl#>
PREFIX rdfs: <http://www.w3.org/2000/01/rdf-schema#>
PREFIX xsd: <http://www.w3.org/2001/XMLSchema#>
PREFIX : <http://www.semanticweb.org/2022/04/aaenote#>
SELECT ?indv_location ?class_catheter
WHERE {
  ?indv_catheter rdf:type/rdfs:subClassOf* :catheter ;
                rdf:type ?class_catheter .
  FILTER(?class_catheter != owl:NamedIndividual) .
  ?indv_location rdf:type/rdfs:subClassOf* :anatomical_location ;
                 rdf:type ?class_location .
  FILTER(?class_location != owl:NamedIndividual) .
  ?indv_catheter :located_nearby_on_at_in ?indv_location .
}
ORDER BY ?indv_location

```

Listing 13: List the anatomical location and the type of catheter located nearby/on/at/in there.

Table 16
AAENOTE SPARQL Query Result: List of patients with a catheter

Query Result	Number of specific catheters	Person Patient	Catheter							
			catheter	central venous catheter	epidural catheter	hickman catheter	hemodialysis catheter	peripheral intravenous catheter	urinary catheter	vas catheter
1	1	patient.T1.1.lo12a.LO_0007						✓		
2	1	patient.T1.1.lo22a.LO_0007						✓		
3	1	patient.T1.1.lo22a.LO_0009					✓			
4	1	patient.T1.2.do13a.LO_0001						✓		
5	1	patient.T1.2.do13a.SP_0001							✓	
6	1	patient.T1.2.do13a.SP_0005						✓		
7	1	patient.T1.2.do13a.SP_0006						✓		
8	1	patient.T1.2.do13a.SP_0010						✓		
9	1	patient.T1.2.lo12a.PO_0005						✓		
10	1	patient.T1.2.lo12a.SD_0007								✓
11	1	patient.T1.2.lo22a.SD_0002						✓		
12	1	patient.T1.2.lo22a.SD_0007								✓
13	1	patient.T1.2.po24a.DO_0001			✓					
...										
22	1	patient.T1.4.lo12a.SP_0003		✓						
23	1	patient.T1.4.lo22a.DO_0003								✓
24	1	patient.T1.4.lo22a.SP_0005						✓		
25	1	patient.T1.4.lo22a.SP_0006						✓		
26	1	patient.T1.4.lo22a.SP_0010						✓		
27	1	patient.T1.4.po24a.LD_0004				✓				
28	1	patient.T1.4.po24a.LD_0009	✓							
29	1	patient.T1.4.po24a.LD_0009		✓						
30	1	patient.T1.4.po24a.LD_0009			✓					
...										
93	1	patient.T3.4.so11a.LO_0003								✓
94	1	patient.T3.4.so21a.LO_0004						✓		
95	2	patient.T4.1.lo22a.LO_0003						✓		
...										
138	1	patient.T9.2.so11a.LD_0003						✓		
139	1	patient.T9.3.po24a.SD_0005						✓		

Query result 1-139: Patient $\xrightarrow{\text{Person has}}$ catheter. And provide the specific type of catheter.
 Query result 28-30, same patient but different catheters.
 Query result 95, the patient has 2 PIVCs.

Table 17
AAENOTE SPARQL Query Result: List anatomical locations with catheters

Query Result	Anatomical Location	Catheter		
	Individuals under class	catheter	venous catheter	peripheral intravenous catheter
1	arm.T14.3.po24a.SD_0002			✓
2	arm.T2.3.po24a.LO_0002			✓
3	arm.T2.3.po24a.SD_0001			✓
4	arm.T3.3.po24a.LO_0005			✓
5	arm.T5.3.po24a.LO_0004			✓
6	arm.T5.3.po24a.SD_0003			✓
7	arm.T5.3.so11a.DO_0002			✓
8	arm.T5.4.lo12a.DO_0003			✓
9	arm.T5.4.lo22a.DO_0003		✓	
	...			
14	arm.T9.4.lo12a.DO_0003			✓
15	body_part.T8.3.po14a.LO_0004			✓
16	elbow.T2.2.lo12a.SD_0001			✓
	...			
69	hand.T9.2.so21a.LD_0003			✓
70	navel.T5.3.so11a.LP_0010	✓		
71	navel.T5.4.do23a.LP_0010	✓		
72	skin.T11.3.po24a.LO_0001			✓
73	skin.T5.2.do23a.LO_0005			✓
74	subcutaneous.T12.2.do23a.LO_0004			✓
75	wrist.T2.2.do13a.LO_0005			✓
76	wrist.T3.2.do23a.LO_0005			✓
77	wrist.T3.4.so11a.LO_0005			✓
78	wrist.T7.3.po14a.LO_0005			✓

Query result 1-78: Catheter ^{Located nearby/on/at/in} anatomical location. And provide the specific anatomical location and type of catheter.

B.10. AAENOTE CQ 7: Does patientB have a PIVC?

This competency question (CQ) can be answered similarly to Appendix B.9. Explicitly using Listing 14 (Table 18) and implicitly using Listing 15 (Table 19). Likewise, additional clinical knowledge is needed to determine if other observations combined indicate a PIVC is present.

```

PREFIX rdf: <http://www.w3.org/1999/02/22-rdf-syntax-ns#>
PREFIX owl: <http://www.w3.org/2002/07/owl#>
PREFIX rdfs: <http://www.w3.org/2000/01/rdf-schema#>
PREFIX xsd: <http://www.w3.org/2001/XMLSchema#>
PREFIX : <http://www.semanticweb.org/2022/04/aaenote#>
SELECT ?indv_patient ?class_pivc (COUNT (DISTINCT ?indv_pivc) as ?n_catheters)
WHERE {
  ?indv_patient rdf:type :patient ;
               rdf:type ?class_patient .
  FILTER(?class_patient != owl:NamedIndividual) .
  ?indv_pivc rdf:type/rdfs:subClassOf* :peripheral_intravenous_catheter ;
            rdf:type ?class_pivc .
  FILTER(?class_pivc != owl:NamedIndividual) .
  ?indv_patient :person_has ?indv_pivc .
}
GROUP BY ?indv_patient ?class_pivc
ORDER BY ?indv_patient
    
```

Listing 14: List patients that have a PIVC and number of PIVCs.

Table 18
AAENOTE SPARQL Query Result: List patients that have a PIVC

Query Result	Number of PIVCs	Patient	peripheral intravenous catheter
1	1	patient.T1.1.lo12a.LO_0007	✓
2	1	patient.T1.1.lo22a.LO_0007	✓
3	1	patient.T1.2.do13a.LO_0001	✓
4	1	patient.T1.2.do13a.SP_0005	✓
5	1	patient.T1.2.do13a.SP_0006	✓
...			
31	1	patient.T2.1.lo22a.LO_0001	✓
32	2	patient.T2.1.lo22a.LO_0003	✓
33	1	patient.T2.2.do23a.LO_0001	✓
...			
91	1	patient.T8.4.lo22a.SP_0010	✓
92	1	patient.T9.2.so11a.LD_0003	✓
93	1	patient.T9.3.po24a.SD_0005	✓

Query result 1-93: Patient $\xrightarrow{\text{Person has}}$ PIVC.

```

1 PREFIX rdf: <http://www.w3.org/1999/02/22-rdf-syntax-ns#>
2 PREFIX owl: <http://www.w3.org/2002/07/owl#>
3 PREFIX rdfs: <http://www.w3.org/2000/01/rdf-schema#>
4 PREFIX xsd: <http://www.w3.org/2001/XMLSchema#>
5 PREFIX : <http://www.semanticweb.org/2022/04/aaenote#>
6 SELECT ?indv_location ?class_pivc
7 WHERE {
8   ?indv_pivc rdf:type/rdfs:subClassOf* :peripheral_intravenous_catheter ;
9     rdf:type ?class_pivc .
10  FILTER(?class_pivc != owl:NamedIndividual) .
11  ?indv_location rdf:type/rdfs:subClassOf* :anatomical_location ;
12     rdf:type ?class_location .
13  FILTER(?class_location != owl:NamedIndividual) .
14  ?indv_pivc :located_nearby_on_at_in ?indv_location .
15 }
16 ORDER BY ?indv_location

```

Listing 15: List anatomical locations with a PIVC.

Table 19
AAENOTE SPARQL Query Result: List anatomical locations with a PIVC

Query Result	Anatomical Location Individuals under class	Catheter peripheral intravenous catheter
1	arm.T14.3.po24a.SD_0002	✓
2	arm.T2.3.po24a.LO_0002	✓
3	arm.T2.3.po24a.SD_0001	✓
4	arm.T3.3.po24a.LO_0005	✓
5	arm.T5.3.po24a.LO_0004	✓
	...	
14	body_part.T8.3.po14a.LO_0004	✓
15	elbow.T2.2.lo12a.SD_0001	✓
16	elbow.T2.2.lo22a.SD_0001	✓
	...	
67	hand.T7.2.lo22a.PO_0005	✓
68	hand.T9.2.so21a.LD_0003	✓
69	skin.T11.3.po24a.LO_0001	✓
70	skin.T5.2.do23a.LO_0005	✓
71	subcutaneous.T12.2.do23a.LO_0004	✓
72	wrist.T2.2.do13a.LO_0005	✓
73	wrist.T3.2.do23a.LO_0005	✓
74	wrist.T3.4.so11a.LO_0005	✓
75	wrist.T7.3.po14a.LO_0005	✓

Query result 1-75: PIVC Located nearby/on/at/in anatomical location. And provide the specific anatomical location.

B.11. AAENOTE CQ 8a: How many catheters does patientB have?

Listing 16 is used to explicitly query the number and types of catheters a patient has as shown in Table 20. Sometimes there is a direct relationship between a person and multiple catheters such as in Query Result 28, 30, 45, and 52 of Table 20. Whereas, typically there is only one direct relationship between one catheter and an anatomical location when using the implicit query Listing 17 (Table 21). Similarly to Appendix B.7 where with clinical knowledge it could be possible to identify infection indications based on a combination of observations at a certain anatomical location, it could also be possible to determine if combinations of observations at certain anatomical locations are indications of a catheter.

```

PREFIX rdf: <http://www.w3.org/1999/02/22-rdf-syntax-ns#>
PREFIX owl: <http://www.w3.org/2002/07/owl#>
PREFIX rdfs: <http://www.w3.org/2000/01/rdf-schema#>
PREFIX xsd: <http://www.w3.org/2001/XMLSchema#>
PREFIX : <http://www.semanticweb.org/2022/04/aaenote#>
# Group ?class_catheters alphabetically by ?indv_patient
SELECT ?indv_patient (GROUP_CONCAT (DISTINCT str(?class_catheter); separator=", " ) as ?catheters)
(COUNT (DISTINCT ?indv_catheter ) as ?n_catheters)
WHERE {
  {
    # Organize alphabetically ?class_catheter
    SELECT ?indv_patient ?indv_catheter ?class_catheter
    WHERE {
      ?indv_patient rdf:type :patient ;
      rdf:type ?class_patient .
      FILTER(?class_patient != owl:NamedIndividual) .
      ?indv_catheter rdf:type/rdfs:subClassOf* :catheter ;
      rdf:type ?class_catheter .
      FILTER(?class_catheter != owl:NamedIndividual) .
      ?indv_patient :person_has ?indv_catheter .
    }
    GROUP BY ?indv_patient ?indv_catheter ?class_catheter
    ORDER BY ?class_catheter
  }
}
GROUP BY ?indv_patient
ORDER BY ?indv_patient

```

Listing 16: Count the number of catheters a patient has and provide the types of catheters.

Table 20

AAENOTE SPARQL Query Result: List how many catheters and type of catheter a patient has

Query Results	Number of catheters	Person	Catheter								
		Patient	catheter	central venous catheter	epidural catheter	hickman catheter	hemodialysis catheter	peripheral intravenous catheter	urinary catheter	vas catheter	venous catheter
1	1	patient.T1.1.lo12a.LO_0007						✓			
2	1	patient.T1.1.lo22a.LO_0007						✓			
3	1	patient.T1.1.lo22a.LO_0009					✓				
4	1	patient.T1.2.do13a.LO_0001						✓			
5	1	patient.T1.2.do13a.SP_0001							✓		
6	1	patient.T1.2.do13a.SP_0005						✓			
7	1	patient.T1.2.do13a.SP_0006						✓			
8	1	patient.T1.2.do13a.SP_0010						✓			
9	1	patient.T1.2.lo12a.PO_0005						✓			
10	1	patient.T1.2.lo12a.SD_0007								✓	
11	1	patient.T1.2.lo22a.SD_0002						✓			
12	1	patient.T1.2.lo22a.SD_0007								✓	
13	1	patient.T1.2.po24a.DO_0001			✓						
			...								
26	1	patient.T1.4.lo22a.SP_0010						✓			
27	1	patient.T1.4.po24a.LD_0004						✓			
28	4	patient.T1.4.po24a.LD_0009	✓	✓	✓				✓		
29	1	patient.T1.4.po24a.SO_0009							✓		
30	2	patient.T1.4.so11a.DP_0002						✓	✓		
			...								
45	2	patient.T2.1.lo22a.LO_0003						✓			
46	1	patient.T2.2.do13a.SP_0003		✓							
47	1	patient.T2.2.do23a.LO_0001						✓			
48	1	patient.T2.2.do23a.LO_0007						✓			
49	1	patient.T2.2.do23a.SP_0006						✓			
50	1	patient.T2.2.lo22a.PO_0005						✓			
51	1	patient.T2.2.po14a.DO_0002						✓			
52	3	patient.T2.2.so11a.LD_0009	✓	✓					✓		
53	1	patient.T2.2.so11a.PO_0005						✓			
			...								
70	1	patient.T3.2.do23a.SP_0003		✓							
71	1	patient.T3.2.po14a.DO_0001			✓						
72	1	patient.T3.2.po14a.DO_0003								✓	
73	1	patient.T3.2.so11a.LD_0003						✓			
74	1	patient.T3.2.so11a.LD_0004			✓						
75	1	patient.T3.3.do13a.SL_0002						✓			
			...								
131	1	patient.T9.2.so11a.LD_0003						✓			
132	1	patient.T9.3.po24a.SD_0005						✓			

Query result 1-132: Patient $\xrightarrow{\text{Person has}}$ catheter. And count how many catheter (s) of a specific type the Patient has.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
511
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

```

PREFIX rdf: <http://www.w3.org/1999/02/22-rdf-syntax-ns#>
PREFIX owl: <http://www.w3.org/2002/07/owl#>
PREFIX rdfs: <http://www.w3.org/2000/01/rdf-schema#>
PREFIX xsd: <http://www.w3.org/2001/XMLSchema#>
PREFIX : <http://www.semanticweb.org/2022/04/aaenote#>
# Group ?class_catheters alphabetically by ?indv_location
SELECT ?indv_location (GROUP_CONCAT (DISTINCT str(?class_catheter); separator=", " ) as ?catheters)
(COUNT (DISTINCT ?indv_catheter ) as ?n_catheters)
WHERE {
  {
    # Organize alphabetically ?class_catheter
    SELECT ?indv_location ?indv_catheter ?class_catheter
    WHERE {
      ?indv_catheter rdf:type/rdfs:subClassOf* :catheter ;
        rdf:type ?class_catheter .
      FILTER(?class_catheter != owl:NamedIndividual) .
      ?indv_location rdf:type/rdfs:subClassOf* :anatomical_location ;
        rdf:type ?class_location .
      FILTER(?class_location != owl:NamedIndividual) .
      ?indv_catheter :located_nearby_on_at_in ?indv_location .
    }
    GROUP BY ?indv_location ?indv_catheter ?class_catheter
    ORDER BY ?class_catheter
  }
}
GROUP BY ?indv_location
ORDER BY ?indv_location

```

Listing 17: Count the number of catheters located nearby/on/at/in an anatomical location and provide the types of catheters.

Table 21
AAENOTE SPARQL Query Result: List how many catheters and type of catheter an anatomical location has

Query Results	Number of catheters	Anatomical Location Individuals under class	Catheter		
			catheter	peripheral intravenous catheter	venous catheter
1	1	arm.T14.3.po24a.SD_0002		✓	
2	1	arm.T2.3.po24a.LO_0002		✓	
3	1	arm.T2.3.po24a.SD_0001		✓	
4	1	arm.T3.3.po24a.LO_0005		✓	
5	1	arm.T5.3.po24a.LO_0004		✓	
6	1	arm.T5.3.po24a.SD_0003		✓	
7	1	arm.T5.3.so11a.DO_0002		✓	
8	1	arm.T5.4.lo12a.DO_0003		✓	
9	1	arm.T5.4.lo22a.DO_0003			✓
		...			
14	1	arm.T9.4.lo12a.DO_0003		✓	
15	1	body_part.T8.3.po14a.LO_0004		✓	
16	1	elbow.T2.2.lo12a.SD_0001		✓	
		...			
67	1	hand.T6.3.do23a.PO_0005		✓	
68	1	hand.T7.2.lo22a.PO_0005		✓	
69	1	hand.T9.2.so21a.LD_0003		✓	
70	1	navel.T5.3.so11a.LP_0010	✓		
71	1	navel.T5.4.do23a.LP_0010	✓		
72	1	skin.T11.3.po24a.LO_0001		✓	
73	1	skin.T5.2.do23a.LO_0005		✓	
74	1	subcutaneous.T12.2.do23a.LO_0004		✓	
75	1	wrist.T2.2.do13a.LO_0005		✓	
76	1	wrist.T3.2.do23a.LO_0005		✓	
77	1	wrist.T3.4.so11a.LO_0005		✓	
78	1	wrist.T7.3.po14a.LO_0005		✓	

Query result 1-78: Catheter $\xrightarrow{\text{Located nearby/on/at/in}}$ anatomical location. And count how many catheter (s) of a specific type the specific anatomical location has.

B.12. AAENOTE CQ 8b: Where are the catheters in patientB?

Explicitly Listing 18 and Table 22. Implicitly, it is the same as Appendix B.9’s Listing 13 and Table 17 where catheter located at an anatomical location indicates a person has the catheter. Here also, additional clinical knowledge is needed to determine if other observations combined indicate a catheter is present.

```

PREFIX rdf: <http://www.w3.org/1999/02/22-rdf-syntax-ns#>
PREFIX owl: <http://www.w3.org/2002/07/owl#>
PREFIX rdfs: <http://www.w3.org/2000/01/rdf-schema#>
PREFIX xsd: <http://www.w3.org/2001/XMLSchema#>
PREFIX : <http://www.semanticweb.org/2022/04/aaenote#>
SELECT ?indv_patient ?indv_catheter ?indv_location
WHERE {
  ?indv_patient rdf:type :patient ;
               rdf:type ?class_patient .
  FILTER(?class_patient != owl:NamedIndividual) .
  ?indv_catheter rdf:type rdfs:subClassOf* :catheter ;
               rdf:type ?class_catheter .
  FILTER(?class_catheter != owl:NamedIndividual) .
  ?indv_location rdf:type rdfs:subClassOf* :anatomical_location ;
               rdf:type ?class_location .
  FILTER(?class_location != owl:NamedIndividual) .
  ?indv_patient :person_has ?indv_catheter ;
               :person_has ?indv_location .
  ?indv_catheter :located_nearby_on_at_in ?indv_location .
}
ORDER BY ?indv_patient
    
```

Listing 18: List patients that have an anatomical location, a catheter, and the patient’s catheter is located near-by/on/at/in the patient’s anatomical location.

Table 22

AAENOTE SPARQL Query Result: Patient has an anatomical location, patient has a catheter, and that catheter is at the patient’s anatomical location.

Query Result	Patient	Catheter	Anatomical Location
1	patient.T1.4.lo22a.DO_0003	venous_catheter.T6.4.lo22a.DO_0003	arm.T5.4.lo22a.DO_0003
2	patient.T4.2.lo22a.SD_0003	peripheral_intravenous_catheter.T5.2.lo22a.SD_0003	hand.T6.2.lo22a.SD_0003

Query result 1-2: Patient (X) $\xrightarrow{\text{Person has}}$ anatomical location (Y), Patient (X) $\xrightarrow{\text{Person has}}$ Catheter (Z), and Catheter (Z) $\xrightarrow{\text{Located nearby/on/at/in}}$ anatomical location (Y).

B.13. AAENOTE CQ 8c: Why does patientB need the catheter(s)?

```

PREFIX rdf: <http://www.w3.org/1999/02/22-rdf-syntax-ns#>
PREFIX owl: <http://www.w3.org/2002/07/owl#>
PREFIX rdfs: <http://www.w3.org/2000/01/rdf-schema#>
PREFIX xsd: <http://www.w3.org/2001/XMLSchema#>
PREFIX : <http://www.semanticweb.org/2022/04/aaenote#>
SELECT ?indv_patient ?class_catheter
      (GROUP_CONCAT (DISTINCT str(?class_procedure); separator=", " ) as ?procedures)
WHERE {
  {
    SELECT ?indv_patient ?class_catheter ?class_procedure
    WHERE {
      ?indv_patient rdf:type :patient ;
        rdf:type ?class_patient .
      FILTER(?class_patient != owl:NamedIndividual) .
      ?indv_catheter rdf:type/rdfs:subClassOf* :catheter ;
        rdf:type ?class_catheter .
      FILTER(?class_catheter != owl:NamedIndividual) .
      ?indv_procedure rdf:type/rdfs:subClassOf* :procedure ;
        rdf:type ?class_procedure .
      FILTER(?class_procedure != owl:NamedIndividual) .
      ?indv_patient :person_has ?indv_catheter .
      ?indv_procedure :procedure_uses ?indv_catheter .
    }
    GROUP BY ?indv_patient ?class_catheter ?class_procedure
    ORDER BY ?class_procedure
  }
}
GROUP BY ?indv_patient ?class_catheter
ORDER BY ?indv_patient

```

Listing 19: List patients that have a catheter and the procedures which use that catheter.

```

PREFIX rdf: <http://www.w3.org/1999/02/22-rdf-syntax-ns#>
PREFIX owl: <http://www.w3.org/2002/07/owl#>
PREFIX rdfs: <http://www.w3.org/2000/01/rdf-schema#>
PREFIX xsd: <http://www.w3.org/2001/XMLSchema#>
PREFIX : <http://www.semanticweb.org/2022/04/aaenote#>
SELECT ?indv_patient ?class_catheter
      (GROUP_CONCAT (DISTINCT str(?class_procedure); separator=", " ) as ?procedures)
WHERE {
  {
    SELECT ?indv_patient ?class_catheter ?class_procedure
    WHERE {
      ?indv_patient rdf:type :patient ;
        rdf:type ?class_patient .
      FILTER(?class_patient != owl:NamedIndividual) .
      ?indv_catheter rdf:type/rdfs:subClassOf* :catheter ;
        rdf:type ?class_catheter .
      FILTER(?class_catheter != owl:NamedIndividual) .
      ?indv_procedure rdf:type/rdfs:subClassOf* :procedure ;
        rdf:type ?class_procedure .
      FILTER(?class_procedure != owl:NamedIndividual) .
      ?indv_patient :person_has ?indv_catheter ;
        :person_has ?indv_procedure .
      ?indv_procedure :procedure_uses ?indv_catheter .
    }
    GROUP BY ?indv_patient ?class_catheter ?class_procedure
    ORDER BY ?class_procedure
  }
}
GROUP BY ?indv_patient ?class_catheter
ORDER BY ?indv_patient

```

Listing 20: List patients that have a catheter, a procedure, and the patient's procedure uses the patient's catheter.

Table 23
AAENOTE SPARQL Query Result: List patients that have a catheter which was used for a procedure

Query Results	Person	Catheter					Procedure									
	Patient	central venous catheter	epidural catheter	peripheral intravenous catheter	venous catheter	urinary catheter	administration purpose	catheter discontinue use	catheter rinse	catheter self removal	general iv	iv antibiotics	iv chemo	iv fluid	iv medication	iv nutrient
1	patient.T1.1.lo12a.LO_0007			✓							✓					
2	patient.T1.1.lo22a.LO_0007			✓							✓		✓			
3	patient.T1.2.do13a.LO_0001			✓							✓					
4	patient.T1.4.lo12a.DO_0003			✓			✓									
5	patient.T1.4.lo12a.SP_0003	✓												✓		
6	patient.T1.4.lo22a.DO_0003				✓		✓									
7	patient.T1.4.lo22a.SP_0010			✓							✓			✓		
8	patient.T1.4.so11a.DP_0002			✓					✓							
9	patient.T1.4.so11a.DP_0002					✓			✓							
10	patient.T1.4.so11a.LO_0001			✓							✓					
11	patient.T11.1.lo22a.LO_0004			✓								✓				
12	patient.T12.4.so11a.LO_0003				✓			✓								
13	patient.T2.2.do23a.LO_0007			✓							✓		✓			
14	patient.T2.3.so11a.LP_0007			✓					✓							
15	patient.T2.3.so21a.LP_0004			✓									✓			
16	patient.T2.4.lo22a.DO_0001	✓					✓									
17	patient.T2.4.so11a.DP_0001			✓											✓	
18	patient.T2.4.so11a.LO_0007			✓							✓					✓
19	patient.T3.2.so11a.LD_0003			✓									✓			
20	patient.T3.4.po24a.SO_0002			✓					✓					✓		
21	patient.T3.4.so11a.LO_0003				✓			✓								
22	patient.T4.3.po14a.SD_0005			✓					✓							
23	patient.T4.3.so11a.LP_0004			✓										✓		
24	patient.T5.3.po24a.LO_0003			✓				✓						✓		
25	patient.T5.4.lo12a.SP_0010			✓							✓					
26	patient.T6.1.lo22a.LO_0007			✓							✓		✓			
27	patient.T6.3.lo22a.DP_0001	✓								✓						
28	patient.T7.2.do23a.LO_0004			✓								✓				
29	patient.T7.3.po24a.SD_0007				✓			✓								
30	patient.T7.4.so11a.LO_0007			✓							✓					✓
31	patient.T8.4.lo22a.SP_0010			✓							✓					
32	patient.T9.3.po24a.SD_0005			✓					✓							

Query result 1-32: Patient $\xrightarrow{\text{Person has}}$ catheter (Y) and Procedure $\xrightarrow{\text{Procedure uses}}$ catheter (Y).

Table 24

AAENOTE SPARQL Query Result: List patients that have a catheter, have a procedure, and where the catheter was used for that procedure

Query Result	Person	Catheter	Procedure		
	Patient	peripheral intravenous catheter	iv antibiotics	iv fluid	iv medication
1	patient.T1.1.lo22a.LO_0007	✓	✓	✓	
2	patient.T1.4.lo22a.SP_0010	✓	✓		✓
3	patient.T2.2.do23a.LO_0007	✓	✓	✓	
4	patient.T6.1.lo22a.LO_0007	✓	✓		✓

Query result 1-14: Patient (X) $\xrightarrow{\text{Person has}}$ catheter (Y), Patient (X) $\xrightarrow{\text{Person has}}$ procedure (Z), and Procedure (Z) $\xrightarrow{\text{Procedure uses}}$ catheter (Y).

```

PREFIX rdf: <http://www.w3.org/1999/02/22-rdf-syntax-ns#>
PREFIX owl: <http://www.w3.org/2002/07/owl#>
PREFIX rdfs: <http://www.w3.org/2000/01/rdf-schema#>
PREFIX xsd: <http://www.w3.org/2001/XMLSchema#>
PREFIX : <http://www.semanticweb.org/2022/04/aaenote#>
SELECT ?indv_patient (GROUP_CONCAT (DISTINCT str(?class_catheter); separator=", " ) as ?catheters)
(GROUP_CONCAT (DISTINCT str(?class_procedure); separator=", " ) as ?procedures)
WHERE {
  {
    SELECT ?indv_patient ?class_catheter ?class_procedure
    WHERE {
      ?indv_patient rdf:type :patient ;
        rdf:type ?class_patient .
      FILTER(?class_patient != owl:NamedIndividual) .
      ?indv_catheter rdf:type/rdfs:subClassOf* :catheter ;
        rdf:type ?class_catheter .
      FILTER(?class_catheter != owl:NamedIndividual) .
      ?indv_procedure rdf:type/rdfs:subClassOf* :procedure ;
        rdf:type ?class_procedure .
      FILTER(?class_procedure != owl:NamedIndividual) .
      ?indv_patient :person_has ?indv_catheter ;
        :person_has ?indv_procedure .
    }
    GROUP BY ?indv_patient ?class_catheter ?class_procedure
    ORDER BY ?class_catheter ?class_procedure
  }
}
GROUP BY ?indv_patient
ORDER BY ?indv_patient

```

Listing 21: List patients that have a catheter and a procedure, and all catheters and all procedures the patient has.

Table 25
AAENOTE SPARQL Query Result: List patients that have a catheter and a procedure

Query Result	Person Patient	Catheter					Procedure							
		catheter	central venous catheter	hemodialysis catheter	peripheral intravenous catheter	urinary catheter	administration purpose	blood test	general iv	iv antibiotics	iv chemo	iv fluid	iv medication	procedure
1	patient.T1.1.lo22a.LO_0007				✓				✓					
2	patient.T1.1.lo22a.LO_0009			✓				✓						
3	patient.T1.2.do13a.SP_0001					✓								✓
4	patient.T1.2.do13a.SP_0010				✓				✓				✓	
5	patient.T1.2.lo22a.SD_0002				✓			✓				✓		
6	patient.T1.4.do13a.LP_0004				✓						✓			
7	patient.T1.4.lo22a.SP_0006				✓			✓						
8	patient.T1.4.lo22a.SP_0010				✓			✓		✓		✓		
9	patient.T12.2.lo22a.SD_0002				✓				✓					
10	patient.T15.3.po24a.SD_0002				✓				✓					
11	patient.T2.1.lo22a.LO_0001				✓				✓					
12	patient.T2.2.do13a.SP_0003		✓									✓		
13	patient.T2.2.do23a.LO_0001				✓				✓					
14	patient.T2.2.do23a.LO_0007				✓				✓		✓			
15	patient.T2.2.lo22a.PO_0005				✓									✓
16	patient.T2.3.do23a.PO_0005				✓									✓
17	patient.T3.2.do13a.LO_0007				✓				✓					
18	patient.T3.2.do23a.SP_0003		✓									✓		
19	patient.T3.3.lo22a.SO_0008	✓										✓		
20	patient.T3.3.po14a.SD_0002				✓							✓		
21	patient.T3.4.lo22a.SP_0003		✓									✓		
22	patient.T4.2.po24a.DO_0002				✓			✓						
23	patient.T5.2.do13a.LO_0003				✓						✓			
24	patient.T5.3.po24a.LO_0001				✓				✓			✓		
25	patient.T6.1.lo22a.LO_0007				✓				✓		✓			
26	patient.T6.2.do13a.LO_0004				✓					✓				
27	patient.T6.4.do13a.LP_0002				✓			✓						
28	patient.T6.4.do23a.LP_0004				✓						✓			
29	patient.T8.4.lo22a.SP_0010				✓			✓						

Query result 1-29: Patient (X) $\xrightarrow{\text{Person has}}$ catheter and Patient (X) $\xrightarrow{\text{Person has}}$ procedure.

B.14. AAENOTE CQ 9a: Does patientC have an infection and catheter?

```

PREFIX rdf: <http://www.w3.org/1999/02/22-rdf-syntax-ns#>
PREFIX owl: <http://www.w3.org/2002/07/owl#>
PREFIX rdfs: <http://www.w3.org/2000/01/rdf-schema#>
PREFIX xsd: <http://www.w3.org/2001/XMLSchema#>
PREFIX : <http://www.semanticweb.org/2022/04/aaenote#>
SELECT ?indv_patient ?class_catheter ?class_infection
WHERE {
  ?indv_patient rdf:type :patient ;
               rdf:type ?class_patient .
  FILTER(?class_patient != owl:NamedIndividual) .
  ?indv_catheter rdf:type/rdfs:subClassOf* :catheter ;
               rdf:type ?class_catheter .
  FILTER(?class_catheter != owl:NamedIndividual) .
  ?indv_infection rdf:type :infection ;
                 rdf:type ?class_infection .
  FILTER(?class_infection != owl:NamedIndividual) .
  ?indv_patient :person_has ?indv_catheter ;
               :person_has ?indv_infection .
}
ORDER BY ?indv_patient
    
```

Listing 22: List patients that have an infection and a catheter.

Table 26
AAENOTE SPARQL Query Result: List patients that have an infection and a catheter

Query Result	Patient	peripheral intravenous catheter	infection
1	patient.T2.2.do23a.SP_0006	✓	✓

Query result 1: Patient (X) $\xrightarrow{\text{Person has}}$ infection and Patient (X) $\xrightarrow{\text{Person has}}$ catheter.

B.15. AAENOTE CQ 9b: Was patientC’s infection associated with a catheter?

Cannot determine if a patient’s infection is associated with a catheter unless that catheter is tested in the microbiology lab.

Appendix C. Catheter Infection Indications Ontology (CIIO) Hierarchy, Assumptions, Indications, and Competency Questions

C.1. CIIO Hierarchy

The CIIO classes and their subclasses can be found in the following figures:

1. Fig. 15 Observation
2. Fig. 16 Anatomical Location
3. Fig. 17 Medical Device
4. Fig. 18 Procedure
5. Fig. 19 Person
6. Fig. 20 Document
7. Fig. 21 Sentence




Fig. 15. CIIO Observation Class Hierarchy




Fig. 16. CIIO Anatomical Location Class Hierarchy




Fig. 17. CIIO Medical Device Class Hierarchy




Fig. 18. CIIO Procedure Class Hierarchy




Fig. 19. CIIO Person Class Hierarchy




Fig. 20. CIIO Document Class Hierarchy




Fig. 21. CIIO Sentence Class Hierarchy

1 C.2. CIO Assumptions 1

2
3 To address the competency questions (CQs) which ask about patients and not documents and alleviate the prob- 3
4 lem of patients being implicitly mentioned, 1 AE document represents 1 patient. In the actual electronic incident 4
5 reporting system database, if the reported AE is related to a patient, there will be a patient ID. This allows users to 5
6 know if the AE is about 1 patient, more than 1 patient, or no patients. Additionally, concepts that are documented 6
7 within different sentences of the same document are likely describing concepts that occurred in the same event for 7
8 the same patient. Furthermore, concepts documented in the same sentence that are linked together by a relationship 8
9 are directly related. Certain relationships can provide the reason for why a concept was or is needed (i.e., procedureA 9
10 Procedure uses → deviceX, therefore deviceX was needed to perform procedureA). 10
11

12 C.3. Catheter and Infection Indications 12

13 C.3.1. Catheter Indications 13

- 14 1. A patient has a specific catheter documented. 14
- 15 2. Any IV usage or infusion indicates some type of catheter is used. IV usage includes general IV, IV medication, 15
16 IV fluid, and IV antibiotics. Infusion includes infusion, intraosseous infusion, intravenous infusion, and subcu- 16
17 taneous infusion. Based on the type of IV usage or infusion alone, it is not enough to determine what type of 17
18 catheter was used. 18
- 19 3. Catheter procedures indicate that a catheter is or was present because they require a catheter. Catheter proce- 19
20 dures include catheter insertion, catheter discontinued use, catheter removal, catheter replacement, and catheter 20
21 self-removal. 21
- 22 4. Infusion phlebitis indication indicates that some type of catheter is used. 22
- 23 23
- 24 24

25 C.3.2. Peripheral Intravenous Catheter (PIVC) Indications 25

- 26 1. PIVCs are rarely documented, so any PIVC explicitly documented indicates a PIVC was used or in use. 26
- 27 2. Leaking IV or an infusion at the arm, elbow, or hand indicates a PIVC is used. IV includes general IV, IV 27
28 medication, IV fluid, and IV antibiotics. Infusion includes infusion, intraosseous infusion, intravenous infusion, 28
29 and subcutaneous infusion. Central venous catheters (CVCs) are deep, so there should not have leakage on the 29
30 skin. PIVC leakage typically occurs because the catheter dressing is not properly secured or the PIVC is placed 30
31 near a movable joint (i.e., elbow) and becomes dislodged. 31
- 32 32

33 C.3.3. Epidural Indication 33

- 34 1. Epidural usage will explicitly be documented. 34
- 35 2. A epidural catheter is the only catheter located nearby/on/at/in the back (i.e., spinal cord). 35
- 36 36

37 C.3.4. Infusion phlebitis 37

38 As previously stated, infusion phlebitis can be mechanical, chemical, or infectious [6]. Regardless of cause, it is 38
39 documented similarly and can either be a catheter-related infection or complication. 39

- 40 1. Early stage of infusion phlebitis is indicated by an insertion site or infusion with 2 of the following signs: (i) 40
41 pain or tenderness, (ii) red, (iii) swollen or edema, or (iv) warm. 41
- 42 2. Medium stage of infusion phlebitis is indicated by (1) a vein with pain and (2) an insertion site or infusion with 42
43 2 signs: (i) red, (ii) swollen or edema, or (iii) warm. 43
- 44 3. Advanced stage of infusion phlebitis is indicated by (1) a vein with pain and hardness and (2) an insertion site 44
45 or infusion with 2 signs: (i) red, (ii) swollen or edema, or (iii) warm. 45
- 46 46

47 C.3.5. Infection 47

- 48 1. Pus at an insertion site indicates an infection. Because pus present is a sure sign of infection. 48
- 49 49

50 C.3.6. Bloodstream Infection (BSI) 50

- 51 1. A bloodstream infection is indicated by a blood test with a positive test result and/or the name of the cultured 51
bacteria. 51

C.3.7. Sepsis

1. Infection combined with mobility impairment, high body temperature, and frostbite indicates sepsis.
2. Meeting the Quick Sequential Organ Failure Assessment Score (qSOFA) [68] sepsis criteria is indicated if there is an infection indication and at least 2 of the following: (1) high respiratory rate, (2) low blood pressure, or (3) a consciousness level that is either confusion, verbally responsive, painfully responsive, or unresponsive.
3. Sepsis is indicated if there is an infection indication and the National Early Warning Score 2 (NEWS2) [69] criteria for clinical deterioration is met by a combination of (1) high respiratory rate, (2) low blood pressure, (3) high pulse, (4) low body temperature or high body temperature, and (5) consciousness level = confusion, verbally responsive, painfully responsive, or unresponsive.

C.4. CIIO Competency Questions and Ontology Usage

SPARQL queries for answering the CIIO competency questions are available on GitHub at https://github.com/melissayan/aaenote_and_cio/wiki/Ontology-SPARQL-Queries.

Table 27: CIIO Competency Questions and Ontology Usage

Competency Question	Ontology Classes and Relationships to Find Instances
1. Does patientA have phlebitis, and was it infectious phlebitis, chemical phlebitis, or mechanical phlebitis?	<p>Early stage infusion phlebitis (a or b):</p> <p>(a) (pain or tenderness), red, (swollen or edema), warm $\xrightarrow{\text{Located nearby/on/at/in}}$ injection site.</p> <p>(b) Infusion $\xrightarrow{\text{Is observed with}}$ (pain or tenderness), red, (swollen or edema), warm.</p> <p>Medium stage infusion phlebitis (a) and (b or c):</p> <p>(a) (pain or tenderness) $\xrightarrow{\text{Located nearby/on/at/in}}$ vein.</p> <p>(b) red, (swollen or edema), warm $\xrightarrow{\text{Located nearby/on/at/in}}$ injection site.</p> <p>(c) Infusion $\xrightarrow{\text{Is observed with}}$ red, (swollen or edema), warm.</p> <p>Advanced stage infusion phlebitis:</p> <p>(a) hardness and (pain or tenderness) $\xrightarrow{\text{Located nearby/on/at/in}}$ vein.</p> <p>(b) red, (swollen or edema), warm $\xrightarrow{\text{Located nearby/on/at/in}}$ injection site.</p> <p>(c) Infusion $\xrightarrow{\text{Is observed with}}$ red, (swollen or edema), warm. [‡]Need additional documentation or pathophysiology studies at the body's cellular or genetic-level to determine if the phlebitis is infectious phlebitis, chemical phlebitis, or mechanical phlebitis.</p>
2. Does patientA have an infection?	<p>Pus at an insertion site indicates infection.</p> <p>Sentence $\xrightarrow{\text{Contains}}$ pus and insertion site. Pus $\xrightarrow{\text{Located nearby/on/at/in}}$ insertion site.</p>
3. Does patientA have a BSI?	<p>Documentation with a blood test with (a) a positive test result and/or (b) name of cultured bacteria.</p> <p>Sentence $\xrightarrow{\text{Contains}}$ blood test.</p> <p>(a) Blood test $\xrightarrow{\text{Has blood test result}}$ Positive.</p> <p>(b) Blood test $\xrightarrow{\text{Has cultured bacteria}}$ bacteria name (i.e., Streptococcus, Staphylococcus, <i>S. aureus</i>, etc.).</p>

1	4. How many patients have an infection or BSI?	Same as Competency Question 1 and 2.	1
2			2
3	5. Which patients have sepsis?	Sepsis is indicated by (a) an infection indication combined with (b): (1) mobility impairment, (2) high body temperature (i.e., hyperthermia), and (3) frostbite:	3
4			4
5		(a) Sentence $\xrightarrow{\text{Contains}}$ pus and insertion site.	5
6		Pus $\xrightarrow{\text{Located nearby/on/at/in}}$ insertion site.	6
7		(b) Sentence $\xrightarrow{\text{Contains}}$ mobility impairment, body	7
8		temperature, and frostbite.	8
9		Body temperature $\xrightarrow{\text{Has body temperature range}}$ hyperthermia.	9
10			10
11		Meeting the Quick Sequential Organ Failure Assessment Score (qSOFA)	11
12		criteria of (a) an infection indication and (b) at least 2 of the following:	12
13		(1) high respiratory rate, (2) low blood pressure, or (3) a consciousness	13
14		level that is either confusion, verbally responsive, painfully responsive, or	14
15		unresponsive:	15
16		(a) Sentence $\xrightarrow{\text{Contains}}$ pus and insertion site.	16
17		Pus $\xrightarrow{\text{Located nearby/on/at/in}}$ insertion site.	17
18		(b) Sentence $\xrightarrow{\text{Contains}}$ respiratory rate, blood	18
19		pressure, and/or consciousness level.	19
20		Respiratory rate $\xrightarrow{\text{Has severity}}$ high.	20
21		Blood pressure $\xrightarrow{\text{Has severity}}$ low.	21
22		Consciousness level $\xrightarrow{\text{Has consciousness state}}$ confusion, ver-	22
23		bally responsive, painfully responsive, or unresponsive.	23
24		Meeting (a) an infection indication and (b) the National Early Warning	24
25		Score 2 (NEWS2) criteria for clinical deterioration by having a combina-	25
26		tion of: (1) high respiratory rate, (2) low blood pressure, (3) high pulse, (4)	26
27		low body temperature or high body temperature (i.e., hypothermia or hy-	27
28		perthermia), and (5) consciousness level = confusion, verbally responsive,	28
29		painfully responsive, or unresponsive:	29
30		(a) Sentence $\xrightarrow{\text{Contains}}$ pus and insertion site.	30
31		Pus $\xrightarrow{\text{Located nearby/on/at/in}}$ insertion site.	31
32		(b) Sentence $\xrightarrow{\text{Contains}}$ respiratory rate, blood	32
33		pressure, pulse, body temperature, and/or	33
34		consciousness level.	34
35		Respiratory rate $\xrightarrow{\text{Has severity}}$ high.	35
36		Blood pressure $\xrightarrow{\text{Has severity}}$ low.	36
37		Pulse $\xrightarrow{\text{Has severity}}$ high.	37
38		Body temperature $\xrightarrow{\text{Has body temperature range}}$ hypothermia or hy-	38
39		perthermia.	39
40		Consciousness level $\xrightarrow{\text{Has consciousness state}}$ confusion, ver-	40
41		bally responsive, painfully responsive, or unresponsive.	41
42			42
43			43
44			44
45			45
46			46
47			47
48			48
49			49
50			50
51			51

6. Does patientB have a catheter?

Documentation of a specific catheter, IV usage, infusion, or catheter-related procedure:

Sentence $\xrightarrow{\text{Contains}}$ medical device(s) or IV-related, infusion-related, or catheter procedure-related procedures.

Early stage infusion phlebitis is indicated by (a) an injection site or (b) infusion with 2 of the following signs: (i) pain or tenderness, (ii) red, (iii) swollen or edema, or (iv) warm:

(a) Sentence $\xrightarrow{\text{Contains}}$ injection site and pain or tenderness, red, swollen or edema, and/or warm.

Pain or tenderness, red, swollen or edema, or warm $\xrightarrow{\text{Located nearby/on/at/in}}$ injection site.

(b) Sentence $\xrightarrow{\text{Contains}}$ infusion and pain or tenderness, red, swollen or edema, and/or warm.

Pain or tenderness, red, swollen or edema, or warm $\xrightarrow{\text{Is observed with}}$ infusion.

Medium stage infusion phlebitis is indicated by (a) a vein with pain or tenderness, and (b) an injection site or (c) infusion with 2 of the following signs: (i) red, (ii) swollen or edema, or (iii) warm:

(a) Sentence $\xrightarrow{\text{Contains}}$ vein and pain or tenderness.

Pain or tenderness $\xrightarrow{\text{Located nearby/on/at/in}}$ vein.

(b) Sentence $\xrightarrow{\text{Contains}}$ injection site and red, swollen or edema, and/or warm.

Red, swollen or edema, or warm $\xrightarrow{\text{Located nearby/on/at/in}}$ injection site.

(c) Sentence $\xrightarrow{\text{Contains}}$ infusion and pain or tenderness, red, swollen or edema, and/or warm.

Red, swollen or edema, or warm $\xrightarrow{\text{Is observed with}}$ infusion.

Advanced stage infusion phlebitis is indicated by (a) a vein with (i) hardness and (ii) pain or tenderness, and (b) an injection site or (c) infusion with 2 of the following signs: (i) red, (ii) swollen or edema, or (iii) warm:

(a) Sentence $\xrightarrow{\text{Contains}}$ vein, hardness, and pain or tenderness.

Hardness and pain or tenderness $\xrightarrow{\text{Located nearby/on/at/in}}$ vein.

(b) Sentence $\xrightarrow{\text{Contains}}$ injection site and red, swollen or edema, and/or warm.

Red, swollen or edema, or warm $\xrightarrow{\text{Located nearby/on/at/in}}$ injection site.

(c) Sentence $\xrightarrow{\text{Contains}}$ infusion and pain or tenderness, red, swollen or edema, and/or warm.

Red, swollen or edema, or warm $\xrightarrow{\text{Is observed with}}$ infusion.

7. Does patientB have a PIVC?

PIVCs are rarely documented, so any PIVC explicitly documented indicates a PIVC was used or in use:

Sentence $\xrightarrow{\text{Contains}}$ PIVC.

Leaking IV or infusion at the arm, elbow or hand indicates PIVC usage:

Sentence $\xrightarrow{\text{Contains}}$ leakage and IV-related or infusion-related procedures.

Leakage $\xrightarrow{\text{Is observed with}}$ IV-related or infusion-related procedures.

1	8a. How many catheters does patientB have?	Same as Competency Question 5’s documentation of a specific catheter, IV usage, infusion, or catheter-related procedure:	1
2			2
3		Sentence $\xrightarrow{\text{Contains}}$ medical device(s) or IV-related,	3
4		infusion-related, or catheter procedure-related procedures.	4
5			5
6		* The exact number of catheters per document cannot be counted because multiple sentences within the document could be describing the same catheter and documented procedures can use the same catheter.	6
7			7
8			8
9	8b. Where are the catheters in patientB?	Medical device located nearby/on/at/in an anatomical location:	9
10			10
11		Sentence $\xrightarrow{\text{Contains}}$ anatomical location and medical device.	11
12		Medical device $\xrightarrow{\text{Located nearby/on/at/in}}$ anatomical location.	12
13			13
14		IV usage, infusion, or catheter-related procedure located nearby/on/at/in an anatomical location:	14
15			15
16		Sentence $\xrightarrow{\text{Contains}}$ anatomical location and IV-related, infusion-related, or catheter procedure-related procedures.	16
17			17
18		IV-related, infusion-related, or catheter procedure-related procedures $\xrightarrow{\text{Located nearby/on/at/in}}$ anatomical location.	18
19			19
20			20
21			21
22		* The exact anatomical location of catheters per document cannot be determined because multiple sentences within a document could be describing the same catheter at the same location but with more general terms (i.e., arm instead of hand), the location’s position was not documented (e.g., If a sentence contains elbowA, right handB, and right armC, then handB is part of armC, but elbowA might or might not be part of armC), multiple procedures can be performed at the same location, and an additional anatomical ontology is needed to infer the location based on catheter type.	22
23			23
24			24
25			25
26			26
27			27
28			28
29			29
30	8c. Why does patientB need the catheter(s)?	The medical device is needed and used in a specific procedure.	30
31			31
32		Sentence $\xrightarrow{\text{Contains}}$ medical device and procedure.	32
33		Procedure $\xrightarrow{\text{Procedure uses}}$ medical device.	33
34		Document has sentences with medical devices and/or procedures:	34
35		Sentence $\xrightarrow{\text{Contains}}$ medical device and/or procedure.	35
36		* The exact reason cannot be determined unless the $\xrightarrow{\text{Procedure uses}}$ object property links procedure and medical device because the indications do not provide a list of reasons for why a specific catheter can be used. However, a clinician can view the retrieved list of devices and procedures to determine if the devices in a document could be used for the procedures documented.	36
37			37
38			38
39			39
40			40
41			41
42	9a. Does patientC have an infection and catheter?	Document with (a) infection indication and (b or c) catheter indication :	42
43			43
44		(a) Sentence $\xrightarrow{\text{Contains}}$ pus and insertion site.	44
45		Pus $\xrightarrow{\text{Located nearby/on/at/in}}$ insertion site.	45
46		(b) Sentence $\xrightarrow{\text{Contains}}$ medical device.	46
47		(c) Sentence $\xrightarrow{\text{Contains}}$ anatomical location and IV-related, infusion-related, or catheter procedure-related procedures.	47
48			48
49	9b. Was patientC’s infection associated with a catheter?	§ Cannot determine if an infection is associated with a catheter unless that catheter is tested in the microbiology lab.	49
50			50
51			51

References

- [1] M.Y. Yan, L.H. Høvik, L.T. Gustad and Ø. Nytrø, Understanding and Reasoning About Early Signs of Sepsis: From Annotation Guideline to Ontology, in: *IEEE International Conference on Bioinformatics and Biomedicine, BIBM 2021, Houston, TX, USA, December 9-12, 2021*, IEEE, 2021, pp. 1906–1911. doi:10.1109/BIBM52615.2021.9669311.
- [2] M.Y. Yan, L.H. Høvik, A. Pedersen, L.T. Gustad and Ø. Nytrø, Preliminary Processing and Analysis of an Adverse Event Dataset for Detecting Sepsis-Related Events, in: *IEEE International Conference on Bioinformatics and Biomedicine, BIBM 2021, Houston, TX, USA, December 9-12, 2021*, IEEE, 2021, pp. 1605–1610. doi:10.1109/BIBM52615.2021.9669410.
- [3] E. Alexandrou, G. Ray-Barruel, P.J. Carr, S.A. Frost, S. Inwood, N.S. Higgins, F. Lin, L. Alberto, L.A. Mermel and C.M. Rickard, Use of Short Peripheral Intravenous Catheters: Characteristics, Management, and Outcomes Worldwide, *Journal of Hospital Medicine* **13** 5 (2018). doi:10.12788/jhm.3039.
- [4] C. Dendle, R.L. Stuart and D. Egerton-Warburton, Think before you insert an intravenous catheter, *The Medical journal of Australia* **199**(10) (2013), 655. doi:10.5694/mja13.10733.
- [5] G. Ray-Barruel, D.F. Polit, J.E. Murfield and C.M. Rickard, Infusion phlebitis assessment measures: a systematic review, *Journal of Evaluation in Clinical Practice* **20**(2) (2014), 191–202. doi:10.1111/jep.12107.
- [6] R. Higginson and A. Parry, Phlebitis: treatment, care and prevention, *Nursing Times* **107**(36) (2011), 18–21.
- [7] L. Zhang, S. Cao, N. Marsh, G. Ray-Barruel, J. Flynn, E. Larsen and C.M. Rickard, Infection risks associated with peripheral vascular catheters, *Journal of Infection Prevention* **17**(5) (2016), 207–213. doi:10.1177/1757177416655472.
- [8] L.E. Huerta and T.W. Rice, Pathologic Difference between Sepsis and Bloodstream Infections, *The Journal of Applied Laboratory Medicine* **3**(4) (2019), 654–663. doi:10.1373/jalm.2018.026245.
- [9] J. Paulsen, A. Mehl, Å. Askim, E. Solligård, B.O. Åsvold and J.K. Damås, Epidemiology and outcome of Staphylococcus aureus bloodstream infection and sepsis in a Norwegian county 1996–2011: an observational study, *BMC Infectious Diseases* **15** (2015), 116. doi:10.1186/s12879-015-0849-4.
- [10] K.E. Rudd, N. Kissoon, D. Limmathurotsakul, S. Bory, B. Mutahunga, C.W. Seymour, D.C. Angus and T.E. West, The global burden of sepsis: barriers and potential solutions, *Critical Care* **22**(1) (2018), 232. doi:10.1186/s13054-018-2157-z.
- [11] L.A. Mermel, Short-term Peripheral Venous Catheter-Related Bloodstream Infections: A Systematic Review, *Clinical Infectious Diseases* **65**(10) (2017), 1757–1762. doi:10.1093/cid/cix562.
- [12] J.M. Rohde, A.J. Odden, C. Bonham, L. Kuhn, P.N. Malani, L.M. Chen, S.A. Flanders and T.J. Iwashyna, The epidemiology of acute organ system dysfunction from severe sepsis outside of the intensive care unit, *Journal of Hospital Medicine* **8**(5) (2013), 243–247. doi:10.1002/jhm.2012.
- [13] C.J. Mungall, C. Torniai, G.V. Gkoutos, S.E. Lewis and M.A. Haendel, Uberon, an integrative multi-species anatomy ontology, *Genome Biology* **13**(1) (2012), R5. doi:10.1186/gb-2012-13-1-r5.
- [14] M. Bada, M. Eckert, M. Palmer and L. Hunter, An Overview of the CRAFT Concept Annotation Guidelines, in: *Proceedings of the Fourth Linguistic Annotation Workshop, LAW 2010, Uppsala, Sweden, July 15-16, 2010*, Association for Computational Linguistics, 2010, pp. 207–211.
- [15] M. Bada, N. Vasilevsky, W.A. Baumgartner, M. Haendel and L.E. Hunter, Gold-standard ontology-based anatomical annotation in the CRAFT Corpus, *Database* **2017** (2017). doi:10.1093/database/bax087.
- [16] B. Smith, W. Ceusters, B. Klagges, J. Köhler, A. Kumar, J. Lomax, C. Mungall, F. Neuhaus, A.L. Rector and C. Rosse, Relations in biomedical ontologies, *Genome Biology* **6**(5) (2005), R46. doi:10.1186/gb-2005-6-5-r46.
- [17] M. Bada and L. Hunter, Desiderata for ontologies to be used in semantic annotation of biomedical documents, *Journal of Biomedical Informatics* **44**(1) (2011), 94–101. doi:10.1016/j.jbi.2010.10.002.
- [18] R.S. Travillian, T. Adamusiak, T. Burdett, M. Gruenberger, J. Hancock, A.-M. Mallon, J. Malone, P. Schofield and H. Parkinson, Anatomy ontologies and potential users: bridging the gap, *Journal of Biomedical Semantics* **2** Suppl 4 (2011), S3. doi:10.1186/2041-1480-2-S4-S3.
- [19] C. Rosse and J.L.V. Mejino Jr, A reference ontology for biomedical informatics: the Foundational Model of Anatomy, *Journal of Biomedical Informatics* **36**(6) (2003), 478–500. doi:10.1016/j.jbi.2003.11.007.
- [20] C. Rosse and J.L.V. Mejino, The Foundational Model of Anatomy Ontology, in: *Anatomy Ontologies for Bioinformatics: Principles and Practice*, A. Burger, D. Davidson and R. Baldock, eds, Springer London, London, 2008, pp. 59–117. ISBN 9781846288852. doi:10.1007/978-1-84628-885-2_4.
- [21] O. Bodenreider, The Unified Medical Language System (UMLS): integrating biomedical terminology, *Nucleic Acids Research* **32**(Database issue) (2004), D267–70. doi:10.1093/nar/gkh061.
- [22] L. Amos, D. Anderson, S. Brody, A. Ripple and B.L. Humphreys, UMLS users and uses: a current overview, *Journal of the American Medical Informatics Association* (2020). doi:10.1093/jamia/ocaa084.
- [23] A.R. Aronson, Effective mapping of biomedical text to the UMLS Metathesaurus: the MetaMap program, *Proceedings. AMIA Symposium* (2001), 17–21.
- [24] V. Ng, Machine Learning for Entity Coreference Resolution: A Retrospective Look at Two Decades of Research, in: *Proceedings of the Thirty-First AAAI Conference on Artificial Intelligence, February 4-9, 2017, San Francisco, California, USA*, Vol. 31, AAAI Press, 2017, pp. 4877–4884. ISSN 2374-3468, 2374-3468. doi:10.1609/aaai.v31i1.11149.
- [25] S. Pradhan, L. Ramshaw, M. Marcus, M. Palmer, R. Weischedel and N. Xue, CoNLL-2011 Shared Task: Modeling Unrestricted Coreference in OntoNotes, in: *Proceedings of the Fifteenth Conference on Computational Natural Language Learning: Shared Task, CoNLL 2011, Portland, Oregon, USA, June 23-24, 2011*, Association for Computational Linguistics, 2011, pp. 1–27.
- [26] S. Pradhan, A. Moschitti, N. Xue, O. Uryupina and Y. Zhang, CoNLL-2012 Shared Task: Modeling Multilingual Unrestricted Coreference in OntoNotes, in: *Joint Conference on Empirical Methods in Natural Language Processing and Computational Natural Language Learning - Proceedings of the Shared Task: Modeling Multilingual Unrestricted Coreference*

- in *OntoNotes*, EMNLP-CoNLL 2012, July 13, 2012, Jeju Island, Korea, Association for Computational Linguistics, 2012, pp. 1–40.
- [27] E. Hovy, M. Marcus, M. Palmer, L. Ramshaw and R. Weischedel, *OntoNotes: The 90% Solution*, in: *Human Language Technology Conference of the North American Chapter of the Association of Computational Linguistics, Proceedings, June 4-9, 2006, New York, New York, USA*, Association for Computational Linguistics, 2006, pp. 57–60.
- [28] S.S. Pradhan, E. Hovy, M. Marcus, M. Palmer, L. Ramshaw and R. Weischedel, *OntoNotes: A Unified Relational Semantic Representation*, in: *Proceedings of the First IEEE International Conference on Semantic Computing (ICSC 2007), September 17-19, 2007, Irvine, California, USA, 2007*, pp. 517–526. doi:10.1109/ICSC.2007.83.
- [29] L.-C. Yu, C.-H. Wu, A. Philpot and E. Hovy, *OntoNotes: Sense Pool Verification Using Google N-gram and Statistical Tests*, in: *Proceedings of the OntoLex Workshop at the 6th International Semantic Web Conference (ISWC 2007), 2007*.
- [30] A. Philpot, E. Hovy and P. Pantel, *The Omega Ontology*, in: *Proceedings of OntoLex 2005 - Ontologies and Lexical Resources, Jeju Island, Korea, October 15, 2005, 2005*.
- [31] M. Uschold and M. King, *Towards a Methodology for Building Ontologies*, in: *Workshop on Basic Ontological Issues in Knowledge Sharing, held in conjunction with IJCAI-95, Montreal, Canada*, Edinburgh: Artificial Intelligence Applications Institute, 1995, pp. 1–13.
- [32] M. Grüninger and M.S. Fox, *Methodology for the Design and Evaluation of Ontologies*, in: *Proceedings of the Workshop on Basic Ontological Issues in Knowledge Sharing, International Joint Conference on Artificial Intelligence (IJCAI-95), Montreal, Canada, 1995*. ISSN 1045-0823.
- [33] M. Fernández-López, A. Gómez-Pérez and N. Juristo, *METHONTOLOGY: From Ontological Art Towards Ontological Engineering*, in: *In Proceedings of the Ontological Engineering AAAI-97 Spring Symposium Series, Stanford, CA, USA, 24–26 March 1997*, AAAI Press, 1997.
- [34] Y. Sure, S. Staab and R. Studer, *On-To-Knowledge Methodology (OTKM)*, in: *Handbook on Ontologies*, S. Staab and R. Studer, eds, Springer Berlin Heidelberg, Berlin, Heidelberg, 2004, pp. 117–132. ISBN 9783540247500. doi:10.1007/978-3-540-24750-0_6.
- [35] M.C. Suárez-Figueroa, A. Gómez-Pérez and M. Fernández-López, *The NeOn Methodology for Ontology Engineering*, in: *Ontology Engineering in a Networked World*, M.C. Suárez-Figueroa, A. Gómez-Pérez, E. Motta and A. Gangemi, eds, Springer Berlin Heidelberg, Berlin, Heidelberg, 2012, pp. 9–34. ISBN 9783642247941. doi:10.1007/978-3-642-24794-1_2.
- [36] O. Corcho, M. Fernández-López and A. Gómez-Pérez, *Methodologies, tools and languages for building ontologies. Where is their meeting point?*, *Data & Knowledge Engineering* **46**(1) (2003), 41–64. doi:10.1016/s0169-023x(02)00195-7.
- [37] H.S. Pinto and J.P. Martins, *Ontologies: How can They be Built?*, *Knowledge and information systems* **6**(4) (2004), 441–464. doi:10.1007/s10115-003-0138-1.
- [38] A. Gómez-Pérez, *Towards a framework to verify knowledge sharing technology*, *Expert Systems with Applications* **11**(4) (1996), 519–529. doi:10.1016/S0957-4174(96)00067-X.
- [39] A. Gómez-Pérez, *Ontology Evaluation*, in: *Handbook on Ontologies*, S. Staab and R. Studer, eds, Springer Berlin Heidelberg, Berlin, Heidelberg, 2004, pp. 251–273. ISBN 9783540247500. doi:10.1007/978-3-540-24750-0_13.
- [40] A. Degbelo, *A snapshot of ontology evaluation criteria and strategies*, in: *Proceedings of the 13th International Conference on Semantic Systems, SEMANTiCS 2017, Amsterdam, The Netherlands, September 11-14, 2017*, ACM, 2017. ISBN 9781450352963. doi:10.1145/3132218.3132219.
- [41] N.F. Noy and C.D. Hafner, *The State of the Art in Ontology Design: A Survey and Comparative Review*, *AI Magazine* **18**(3) (1997), 53–53. doi:10.1609/aimag.v18i3.1306.
- [42] *Infectious Disease Ontology*, 2017. <https://bioportal.bioontology.org/ontologies/IDO>.
- [43] *Ontology for General Medical Science*, 2021. <http://www.obofoundry.org/ontology/ogms>.
- [44] *Vital Sign Ontology*, 2012. <https://bioportal.bioontology.org/ontologies/VSO>.
- [45] A. Goldfain, B. Smith, S. Arabandi, M. Brochhausen and W.R. Hogan, *Vital Sign Ontology*, in: *Proceedings of the Workshop on Bio-Ontologies, Intelligent Systems in Molecular Biology (ISMB), July 17-19, 2011, Vienna, Austria, 2011*.
- [46] *Foundational Model of Anatomy*, 2008. <http://si.washington.edu/projects/fma>.
- [47] *Biological Spatial Ontology*, 2020. <http://obofoundry.org/ontology/bsp>.
- [48] *Ontology of Adverse Events*, 2021. <http://www.obofoundry.org/ontology/oa>.
- [49] Y. He, S. Sarntivijai, Y. Lin, Z. Xiang, A. Guo, S. Zhang, D. Jagannathan, L. Toldo, C. Tao and B. Smith, *OAE: The Ontology of Adverse Events*, *Journal of Biomedical Semantics* **5** (2014), 29. doi:10.1186/2041-1480-5-29.
- [50] *Relation Ontology*, 2021. <http://obofoundry.org/ontology/ro>.
- [51] *National Cancer Institute Thesaurus*. <https://bioportal.bioontology.org/ontologies/NCIT>.
- [52] *NCI Thesaurus*. <https://ncithesaurus.nci.nih.gov/ncitbrowser/>.
- [53] *eHealth & International Classification for Nursing Practice (ICNP™)*, 2021. <https://www.icn.ch/what-we-do/projects/ehealth-icnptm>.
- [54] H.K. Butcher, G.M. Bulechek, J.M. Dochterman and C.M. Wagner, *Nursing Interventions Classification (NIC)*, Elsevier, 2018. ISBN 9780323497701.
- [55] T. Heather Herdman, S. Kamitsuru and C.T. Lopes, *NANDA International Nursing Diagnoses: Definitions & Classification, 2021-2023*, Thieme Medical Publishers, Incorporated, 2021. ISBN 9781684204540.
- [56] A. Jackson, *Infection control—a battle in vein: infusion phlebitis*, *Nursing Times* **94**(4) (1998), 68–71.
- [57] *Infusion Nurses Society, 2021 Infusion Therapy Standards of Practice Updates*, *Journal of Infusion Nursing* **44**(4) (2021), 189–190. doi:10.1097/NAN.0000000000000396.
- [58] P. Stenetorp, S. Pyysalo, G. Topić, T. Ohta, S. Ananiadou and J. Tsujii, *BRAT: a Web-based Tool for NLP-assisted Text Annotation*, in: *EACL 2012, 13th Conference of the European Chapter of the Association for Computational Linguistics, Avignon, France, April 23-27, 2012*, 2012, pp. 102–107.
- [59] R. Salomão, B.L. Ferreira, M.C. Salomão, S.S. Santos, L.C.P. Azevedo and M.K.C. Brunialti, *Sepsis: evolving concepts and challenges*, *Brazilian Journal of Medical and Biological Research* **52**(4) (2019), e8595. doi:10.1590/1414-431X20198595.

- [60] K.N. Iskander, M.F. Osuchowski, D.J. Stearns-Kurosawa, S. Kurosawa, D. Stepien, C. Valentine and D.G. Remick, Sepsis: multiple abnormalities, heterogeneous responses, and evolving understanding, *Physiological Reviews* **93**(3) (2013), 1247–1288. doi:10.1152/physrev.00037.2012.
- [61] J. Kolikof, K. Peterson and A.M. Baker, Central Venous Catheter, in: *StatPearls*, StatPearls Publishing, Treasure Island (FL), 2021.
- [62] H.J. White and M.P. Soos, Anatomy, Thorax, Superior Vena Cava, in: *StatPearls*, StatPearls Publishing, Treasure Island (FL), 2021.
- [63] S.M. Meystre, G.K. Savova, K.C. Kipper-Schuler and J.F. Hurdle, Extracting information from textual documents in the electronic health record: a review of recent research, *Yearbook of Medical Informatics* (2008), 128–144. doi:10.1055/s-0038-1638592.
- [64] L.H. Høvik, K.H. Gjeilo, S. Lydersen, C.M. Rickard, B. Røtvold, J.K. Damås, E. Solligård and L.T. Gustad, Monitoring quality of care for peripheral intravenous catheters; feasibility and reliability of the peripheral intravenous catheters mini questionnaire (PIVC-miniQ), *BMC Health Services Research* **19**(1) (2019), 636. doi:10.1186/s12913-019-4497-z.
- [65] M. Villegas, A. Intxaurreondo, A. Gonzalez-Agirre, M. Marimon and M. Krallinger, The MeSpEN resource for English-Spanish medical machine translation and terminologies: Census of parallel corpora, glossaries and term translations, in: *Proceedings of the LREC 2018 Workshop “MultilingualBIO: Multilingual Biomedical Text Processing”*, European Language Resources Association (ELRA), 2018.
- [66] D. Garijo, WIDOCO: a wizard for documenting ontologies, in: *The Semantic Web - ISWC 2017 - 16th International Semantic Web Conference, Vienna, Austria, October 21-25, 2017, Proceedings, Part II*, Springer, 2017, pp. 94–102. doi:10.1007/978-3-319-68204-4_9.
- [67] M. Sabou and M. Fernandez, Ontology (Network) Evaluation, in: *Ontology Engineering in a Networked World*, M.C. Suárez-Figueroa, A. Gómez-Pérez, E. Motta and A. Gangemi, eds, Springer Berlin Heidelberg, Berlin, Heidelberg, 2012, pp. 193–212. ISBN 9783642247941. doi:10.1007/978-3-642-24794-1_9.
- [68] C.W. Seymour, V.X. Liu, T.J. Iwashyna, F.M. Brunkhorst, T.D. Rea, A. Scherag, G. Rubinfeld, J.M. Kahn, M. Shankar-Hari, M. Singer, C.S. Deutschman, G.J. Escobar and D.C. Angus, Assessment of Clinical Criteria for Sepsis: For the Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3), *JAMA* **315**(8) (2016), 762–774. doi:10.1001/jama.2016.0288.
- [69] Royal College of Physicians, National Early Warning Score (NEWS) 2: Standardising the assessment of acute-illness severity in the NHS. Updated report of a working party, Technical Report, Royal College of Physicians, 2017.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51